ANESTHESIA & ANALGESIA

Journal of the International Anesthesia Research Society, the Society of Cardiovascular Anesthesiologists, the Society for Pediatric Anesthesia, and the Society for Ambulatory Anesthesia

Index
Volume 86, 1998

EDITORIAL BOARD
Anesthesia & Analgesia

Editor-in-Chief
Ronald D. Miller, MD

Associate Editor-in-Chief Cardiovascular Anesthesia
Kenneth J. Tuman, MD

Editors
David J. Birnbach, MD, *Obstetric Anesthesia*
Thomas J. J. Blanck, MD, PhD
Daniel B. Carr, MD
Pierre Coriat, MD
Peter G. Duncan, MD, *Economics and Health Systems Research*
Thomas J. Gal, MD
Simon Gelman, MD, PhD
William J. Greeley, MD, MBA, *Pediatric Anesthesia*
Sten Lindahl, MD
Michael F. Roizen, MD
John C. Rowlingson, MD
Koh Shingu, MD
Graham Smith, BSc, MD, FRCA
Christoph Stein, MD
Peter M. Suter, MD, *Critical Care and Trauma*
Hugo Van Aken, MD, PhD
David S. Warner, MD, *Neurosurgical Anesthesia*
Denise J. Wedel, MD, *Regional Anesthesia and Pain Management*
Paul F. White, MD, PhD, *Ambulatory Anesthesia*

Book and Multimedia Review Editor
Norig Ellison, MD

Statistical Editor
Nathan L. Pace, MD

Author Index

Abdelmalak B, see Sprung J
Abe K, Mashimo T, Yoshiya I. Arterial oxygenation and shunt fraction during one-lung ventilation: a comparison of isoflurane and sevoflurane, 1266
Takada K, Yoshiya I. Intraoperative torsade de pointes ventricular tachycardia and ventricular fibrillation during sevoflurane anesthesia, 701
Abel M, see Bronheim D
Abiko Y, see Kokita N
Abisseror M, see Gueugniaud P-Y
Abouleish AE, Mayhew JF. Magnetic resonance imaging of the airway in an infant with micrognathia, 964
Abouleish EI, see Campos CJ
Acimovic S, see Kettner SC
Adachi T, see Kito K
Adelsmayr E, Keller C, Erd G, Brimacombe J. The laryngeal mask and high-frequency jet ventilation for resection of high tracheal stenosis, 907
Adriaensen HA, see Vercauteren MP
Adsan H, Tulunay M, Onaran O. The effects of verapamil and nimodipine on bupivacaine-induced cardiotoxicity in rats: an *in vivo* and *in vitro* study, 818
Adsumelli RSN, see Steinberg ES
Ahn EK, Bai SJ, Cho BJ, Shin Y-S. The infusion rate of mivacurium and its spontaneous neuromuscular recovery in magnesium-treated parturients, 523
Ajimal GS, see Purdy EP
Akamatsu H, see Nishina K
Aken HV, see Brodner G
Albrechtsen M, see Rosenow DE
Ali-Melkkilä T, see Ranta SO-V
Allen GC, Brubaker CL. Human malignant hyperthermia associated with desflurane anesthesia, 1328
Allen RW, see Robbins WR
Allvin RA, see Rawal N
Alon E, Buchser E, Herrera E, Christiaens F, De Pauw C, Ritter L, Hulstaert F, Grimaudo V. Tropisetron for treating established postoperative nausea and vomiting: a randomized, double-blind, placebo-controlled study, 617
Alston RP, see Souter MJ
Alted-Lopez E, see Sanchez-Izquierdo-Riera JA
Amaro AR, see Cardoso MAMSC
Ambesh SP, Kaushik S. Blind orotracheal intubation with the Augustine Guide: a prospective study, 435
Ambros-Checa A, see Sanchez-Izquierdo-Riera JA
Amilon CA, see Rawal N
An G, see Xue FS
Anderson GD, see Ramamoorthy C
Andrews PJD
see McKeating EG
see Souter MJ
Angelillo IF, see Ballantyne JC
Angst MS. Dose-response function of epidural fentanyl versus sufentanil (letter), 918
Antoun S, see Sicsic J-C
Aono M
see Ide T
see Sakurai Y
Aoyama K, Yamamoto T, Takenaka I, Sata T, Shigematsu A. The jaw support device facilitates laryngeal exposure and ventilation during fiberoptic intubation, 432
Apsner R, see Muhm M
Arakawa J, see Kokita N
Areewatana S, see Sanansilp V
Arendt-Nielsen L
see Curatolo M
see Wilder-Smith OHG
Arndt JO, see Holthusen H
Artru AA, see Talmor D
Asskali F, see Behne M
Atlee JL, see Mahla E
Avram MJ, see Dsida RM
Axelsson K, see Rawal N
Aya AGM, Mangin R, Lalourcey L, Eledjam J-J. Multiport versus uniport epidural catheters (letter), 676

Babb JD, see Owen MD
Bacher A, see Kwon J-Y
Bacon MF, see Nader-Djalal N
Bahk J-H, Oh Y-S. A new and simple maneuver to position the left-sided double-lumen tube without the aid of fiberoptic bronchoscopy, 1271
Bai SJ, see Ahn EK
Baisden CE, see Rathmell JP
Ballantyne JC, Carr DB, deFerranti S, Suarez T, Lau J, Chalmers TC, Angelilio IF, Mosteller F. The comparative effects of postoperative analgesic therapies on pulmonary outcome: cumulative meta-analyses of randomized, controlled trials, 598
Chang YC. Does the choice of muscle relaxants really affect postoperative recovery time? (letter, reply), 1335
Banoub MF, Firestone L, Sprung J. Anesthetic management of a patient undergoing minimally invasive myocardial revascularization before lung transplantation, 939
Bardoczky GI, d'Hollander AA, Cappello M, Yernault J-C. Interrupted expiratory flow on automatically constructed flow-volume curves may determine the presence of intrinsic positive end-expiratory pressure during one-lung ventilation, 880
Barthe A, see Vizquel L
Barthelet Y, see Capdevila X
Barton B, see Katz JJ
Bastani B, see Joh J
Bautista Davis EC. Acute and Chronic Pain, Vol. 25, No. 2 of International Anesthesiology Clinics (book review), 682
Beebe DS, see Koehntop DE
Beers RA, O'Leary CE, Franklin PD. Comparing the history-taking methods used during a preanesthesia visit: the HealthQuiz versus the written questionnaire, 134
Behne M, Thomas H, Bremerich DH, Lischke V, Asskali F, Förster H. The pharmacokinetics of acetyl starch as a plasma volume expander in patients undergoing elective surgery, 856
Behrens U, see Striebel HW
Beilin Y, Bernstein HH, Zucker-Pinchoff B, Zahn J, Zenzen WJ. Subhypnotic doses of propofol do not relieve pruritus induced by intrathecal morphine after cesarean section, 310
Belani KG, see Koehntop DE
Bellomo G, see Imberti R
Benani MR, see Vizquel L
Ben-David B. Intrathecal fentanyl with small-dose dilute bupivacaine: better anesthesia without prolonging recovery (letter, reply), 918
Benedek GA, see Klimscha W

Benhamou D
see Bouaziz H
see Bouaziz HA
see Lentschener C
see Mercier FAAJ
Bennett B, see Lobato EB
Bennett GJ, see Sethna NF
Bennett-Guerrero E, Mythen MG. Could we know ϵ-aminocaproic acid level during cardiopulmonary bypass? (letter, reply), 681
Schwinn DA. Treatment of myxedema coma for emergency surgery (letter, reply), 451
Benter IF, see Zagvazdin Y
Bentzon MW, see Michelsen I
Berger J
see Mahla E
see Mächler HE
Bergmann P, see Trantina AE
Berkowitz RJ, see Litman RS
Berman MF, Simon AE. The effect of a drug and supply cost feedback system on the use of intraoperative resources by anesthesiologists, 510
Bernstein HH, see Beilin Y
Berry AM, see Brimacombe JR
Bertrand MA, see Eyraud D
Bhambani R, see Sprung J
Bhargava M, Pothula SNM. Improvement of pulse oximetry signal by EMLA cream (letter), 915
Biboulet P, see Capdevila X
Bilir BM, see Katz JJ
Billett C, see Kost-Byerly S
Birmingham PK, see Dsida RM
Birnbaum JA, see Haisjackl M
Bischoff P, Mahlstedt D, Blanc I, Schulte am Esch J. Quantitative topographical electroencephalographic analysis after intravenous clonidine in healthy male volunteers, 202
Bishop MJ. Is the Combitube a useful emergency airway device for anesthesiologists?, 1141
Bjoraker R, see Haws JL
Blaicher A, see Kettner SC
Blaicher AM, Reiter WJ, Blaicher W, Kettner SC, Felfernig M, Grabner CM, Zimpfer M. The effect of hydroxyethyl starch on platelet aggregation *in vitro*, 1318
Blaicher W, see Blaicher AM
Blanc I, see Bischoff P
Blanck TJJ, see Nikonorov IM
Blomqvist H. Anesthesia and Uncommon Diseases. 4th ed. (book review), 1151
Bodian C, see Bronheim D
Bogetz MS, Swisher JL. Formal training in anesthesia (letter), 219
Boldt J, Jaun N, Kumle B, Heck M, Mund K. Economic considerations of the use of new anesthetics: a comparison of propofol, sevoflurane, desflurane, and isoflurane, 504
Bone HG, see Meyer JA
Booke M, Van Aken H, Theilmeier G, Meyer JA. Prostaglandins in patients with pulmonary hypertension: the route of administration (letter), 917
see Meyer JA
Borgeat A, Ruetsch YA. Anesthesia in a patient with malignant systemic mastocytosis using a total intravenous anesthetic technique, 442
Bornheim LM, see Yost CS
Boschin M, see Buerkle H
Bothner U, Georgieff M, Vogt NH. Assessment of the safety and tolerance of 6% hydroxyethyl starch (200/0.5) solution: a randomized, controlled epidemiology study, 850
Bouaziz H, Narchi P, Mercier FJ, Khoury A, Poirier T, Benhamou D. The use of a selective axillary nerve block for outpatient hand surgery, 746
Bouaziz HA, Benhamou D. The "midhumeral" approach to the brachial plexus (letter, reply), 448
see Mercier FAAJ
Boulez JL, see Branche PE
Bouregba M, see Capdevila X
Bourzoufi K, see Fleyfel M
Bowland T, see Eger EI II
Brain AIJ, see Brimacombe JR
Branche PE, Duperret SL, Sagnard PE, Boulez JL, Petit PL, Viale J-P. Left ventricular loading modifications induced by pneumoperitoneum: a time course echocardiographic study, 482
Brandom BW, see Simhi E
Brands E, see Sher MH
Bratton SL, see Lynn A
Bremerich DH, see Behne M
Brimacombe JR. Laryngeal mask insertion techniques (letter), 1337
Brain AIJ, Berry AM, Verghese C, Ferson D. Gastric insufflation and the laryngeal mask (letter), 914
see Adelsmayr E
see Keller C
Brinkmann H, see Klimscha W
Brodner G, Pogatzki E, Aken HV, Buerkle H, Goeters C, Schulzki C, Nottberg H, Mertes N. A multimodal approach to control postoperative pathophysiology and rehabilitation in patients undergoing abdominothoracic esophagectomy, 228
see Buerkle H
see Hönemann CW
Brodsky JB, see Coppa GP
Bronheim D, Abel M, Bodian C. Does the choice of muscle relaxants really affect postoperative recovery time? (letter), 1335
Brown B, see Macario A
Brown MM, see Lobato EB
Bruandet N, Rentero N, Debeer L, Quintin L. Catecholamine activation in the vasomotor center on emergence from anesthesia: the effects of α_2 agonists, 240
Brubaker CL, see Allen GC
Bruder N, Cohen B, Pellissier D, François G. The effect of hemodilution on cerebral blood flow velocity in anesthetized patients, 320
Brull SJ, see Kelly DJ
Bryson GL, Laupacis A, Wells GA, International Study of Perioperative Transfusion. Does acute normovolemic hemodilution reduce perioperative allogeneic transfusion? A meta-analysis, 9
Buchser E, see Alon E
Buerkle H, Boschin M, Marcus MAE, Brodner G, Wüsten R, Van Aken H. Central and peripheral analgesia mediated by the acetylcholinesterase-inhibitor neostigmine in the rat inflamed knee joint model, 1027
see Brodner G
see Marcus MAE
Buratti T, see Manfreda SE
Burm AGL
see Groen K
see van den Nieuwenhuyzen MCO
Busse H, see Meyer JA
Butterworth JF, see Rathmell JP
Butterworth JF IV, Legault C, Royster RL, Hammon JW Jr. Factors that predict the use of positive inotropic drug support after cardiac valve surgery, 461
Bynon S, see Frenette L

Caballero-Cubedo RE, see Sanchez-Izquierdo-Riera JA
Campbell JN, see Robbins WR

Campos CJ, Pivalizza EG, Abouleish EI. Thromboelastography in a parturient with immune thrombocytopenic purpura (letter), 675
Campos JH, Massa FC. Is there a better right-sided tube for one-lung ventilation? A comparison of the right-sided double-lumen tube with the single-lumen tube with right-sided enclosed bronchial blocker, 696
Canessa R, see Lema G
Cantalapiedra-Santiago JA, see Sanchez-Izquierdo-Riera JA
Capdevila X, Biboulet P, Bouregba M, Barthelet Y, Rubenovitch J, d'Athis F. Comparison of the three-in-one and fascia iliaca compartment blocks in adults: clinical and radiographic analysis, 1039
Biboulet P, Rubenovitch J, Serre-Cousine O, Peray P, Deschodt J, d'Athis FA. The effects of cervical epidural anesthesia with bupivacaine on pulmonary function in conscious patients, 1033
Caplan AB, see DeLoach LJ
Cappelli EL, see Cardoso MAMSC
Cappello M, see Bardoczky GI
Cardoso MAMSC, Carvalho JACA, Amaro AR, Prado AA, Cappelli EL. Small doses of intrathecal morphine combined with systemic diclofenac for postoperative pain control after cesarean delivery, 538
Carlier M, see De Kock M
Carlisle AS. The ICU Book. 2nd ed. (book review), 1151
Carlsén H-O, see Walldén J
Carmody JJ, see Walker JS
Carr D, see Guptill W
Carr DB, see Ballantyne JC
Carvalho JACA, see Cardoso MAMSC
Cassuto J, see Tarnow P
Castelli I, see Drop L
Celebioglu B. A case of suspected malignant hyperthermia during desflurane administration (letter), 916
Could we know ϵ-aminocaproic acid level during cardiopulmonary bypass? (letter), 680
Chakrabarti MK, see Ma D
Chalmers TC, see Ballantyne JC
Chan BCB, see Lui P-W
Chan FHY, see Lui P-W
Chan MTV, see Gin T
Chan YW, see Chow MYH
Chang C-H, see Tan PPC
Chang H-C, see Chia Y-Y
Chang YC, see Ballantyne JC
Chen YCJ, Gajraj NM, Clavo A, Joshi GP. Posterior subcapsular cataract formation associated with multiple lumbar epidural corticosteroid injections, 1054
Cheng K-S, see Tan PPC
Chhabra B, see Gupta R
Chi OZ, Chun TW, Liu X, Weiss HR. The effects of pentobarbital on blood-brain barrier disruption caused by intracarotid injection of hyperosmolar mannitol in rats, 1230
Chia Y-Y, Liu K, Liu Y-C, Chang H-C, Wong C-S. Adding ketamine in a multimodal patient-controlled epidural regimen reduces postoperative pain and analgesic consumption, 1245
Chiari A, see Klimscha W
Chikusa H, see Tsujimoto S
Cho BJ, see Ahn EK
Choe H, Kim J-S, Ko S-H, Kim D-C, Han Y-J, Song H-S. Epidural verapamil reduces analgesic consumption after lower abdominal surgery, 786
Chou T-C, Li C-Y, Wu C-C, Yen M-H, Ding Y-A. The inhibition by dantrolene of L-arginine transport and nitric oxide synthase in rat alveolar macrophages, 1065
Chow MYH, Sia ATH, Koay CK, Chan YW. Alkalinization of lidocaine does not hasten the onset of axillary brachial plexus block, 566
Christiaens F, see Alon E
Chu J-J, see Tan PPC
Chui PT, see Gin T
Chun TW, see Chi OZ
Chung F, see Tong D
Cicutti NJ, see Rosaeg OP
Clanachan AS, see Lazar ER
Clark Y, see Haws JL
Clavo A, see Chen YCJ
Clergue FA, see Klopfenstein CE
Cnaan A, see Cook-Sather SD
Cohen B, see Bruder N
Cohen MA, see Reuben SS
Cohen MM, see Devitt JH
Cole DJ, see Patel PM
Coleman D, see Zhang J
Colonna-Romano P, Nagaraj L. Tests to evaluate intravenous placement of epidural catheters in laboring women: a prospective clinical study, 985
Colson P. Hemodynamic effects of anesthesia in patients receiving angiotensin-converting enzyme inhibitors (letter, reply), 216
Contreras MG, see Warner ME
Cook-Sather SD, Tulloch HV, Cnaan A, Nicolson SC, Cubina ML, Gallagher PR, Schreiner MS. A comparison of awake versus paralyzed tracheal intubation for infants with pyloric stenosis, 945
Coppa GP, Brodsky JB. A simple method to protect the tracheal cuff of a double-lumen tube (letter), 675
Coppejans HC, see Vercauteren MP
Corbineau HA, see Sicsic J-C
Coriat P, see Eyraud D
Cormack JR, see Costello TG
Correia R, see Sprung J
Costello TG, Cormack JR. Clonidine premedication decreases hemodynamic responses to pin head-holder application during craniotomy, 1001
Coté CJ
see Dsida RM
see Eckmann DM
Cousins MJ, see Jaquenod M
Coveler LA, see Norman PH
Cox J, see Frenette L
Creed MR, see Fortney JT
Croughwell ND, see Newman MF
Csavoy AN, see Segal S
Cubina ML, see Cook-Sather SD
Curatolo M, Petersen-Felix S, Arendt-Nielsen L, Lauber R, Högström H, Scaramozzino P, Luginbühl M, Sieber TJ, Zbinden AM. Adding sodium bicarbonate to lidocaine enhances the depth of epidural blockade, 341
see Sieber TJ
Cuthbertson BH, Galley HF, Webster NR. The effects of nitric oxide and peroxynitrite on interleukin-8 and elastase from lipopolysaccharide-stimulated whole blood, 427
Czech T, see Schramm WM

Dacar D, see Trantina AE
Dagnino JA, see Muñoz HAR
Dai HB, see Park KW
Dalamangas LC, Slaughter TF. Ethylenediaminetetraacetic acid-dependent pseudothrombocytopenia in a cardiac surgical patient, 1210
Daley MD, see Norman PH
Dan K, see Kubo N
Datta S
see Levin A
see Segal S

Davis DA. Pediatric Cardiac Anesthesia (book review), 1341
Davis JD, see Lobato EB
De Kock M, Laterre P-FA, Van Obbergh L, Carlier M, Lerut J. The effects of intraoperative intravenous clonidine on fluid requirements, hemodynamic variables, and support during liver transplantation: a prospective, randomized study, 468
De Pauw C, see Alon E
de Rosayro M, see Mullin V
Debeer L, see Bruandet N
deFerranti S, see Ballantyne JC
DeLoach LJ, Higgins MS, Caplan AB, Stiff JL. The visual analog scale in the immediate postoperative period: intrasubject variability and correlation with a numeric scale, 102
Derrick JL, see Gin T
Deschodt J, see Capdevila X
Desmonts J-M, see Molliex S
Devitt JH, Kurrek MM, Cohen MM, Fish K, Fish P, Noel AG, Szalai J-P. Testing internal consistency and construct validity during evaluation of performance in a patient simulator, 1160
Dexter F, see Macario A
Dieterich H-JA, Kraft D, Sirtl C, Laubenthal H, Schimetta W, Pölz W, Gerlach E, Peter K. Hydroxyethyl starch antibodies in humans: incidence and clinical relevance, 1123
Dietrich W, see Sitzwohl C
DiFazio CA, see Lee VC
Ding Y-A, see Chou T-C
Dobos IA, see Klimscha W
Dodat H, see Gueugniaud P-Y
Dohi S
see Ishiyama T
see Ohata H
Dooley JW, see Richardson MG
Doyle MT, Twomey CF, Owens TM, McShane AJ. Gastroesophageal reflux and tracheal contamination during laparoscopic cholecystectomy and diagnostic gynecological laparoscopy, 624
Driver RP Jr, Snyder IS, North FP, Fife TJ. Sterility of anesthetic and resuscitative drug syringes used in the obstetric operating room, 994
Drop L, Castelli I, Kaufmann M. Comparative doses and cost: esmolol versus labetalol during electroconvulsive therapy (letter), 916
Drummond JC, see Patel PM
Dsida RM, Wheeler M, Birmingham PK, Henthorn TK, Avram MJ, Enders-Klein C, Maddalozzo J, Coté CJ. Premedication of pediatric tonsillectomy patients with oral transmucosal fentanyl citrate, 66
Dubbels AM, see Galley HF
Dubreuil M, see Grenier B
Dugas MJ, see Norman PH
Duncan B, see Fortney JT
Duncan PG. That was then, this is now! The value of observing change (editorial), 225
Dunzendorfer S, see Manfreda SE
Duperret SL, see Branche PE
Duranteau J, see Sicsic J-C
Dureuil B, see Molliex S
Durieux ME, see Hönemann CW
Dzwonczyk RR, see Harter RL
D'Angelo R. Multiport versus uniport epidural catheters (letter, reply), 676
see Owen MD
d'Athis F, see Capdevila X
d'Hollander AA, see Bardoczky GI

Eaton MP. Intrathecal sufentanil analgesia for extracorporeal shock wave lithotripsy in three patients with aortic stenosis, 943
Eberle B
see Latorre F
see Weiler N
Ebert TJ, Messana LD, Uhrich TD, Staacke TS. Absence of renal and hepatic toxicity after four hours of 1.25 minimum alveolar anesthetic concentration sevoflurane anesthesia in volunteers, 662
Eckhoff D, see Frenette L
Eckmann DM, Seligman I, Coté CJ, Hussong JW. Mucormycosis supraglottitis on induction of anesthesia in an immunocompromised host, 729
Eckstein RP, see Jaquenod M
Ecoffey C, see Sicsic J-C
Ederberg S, Westerlind A, Houltz E, Svensson S-E, Elam M, Ricksten S-E. The effects of propofol on cerebral blood flow velocity and cerebral oxygen extraction during cardiopulmonary bypass, 1201
Eger EI II. Ethics of human toxicity study questioned (letter, reply), 445
Gong D, Koblin DD, Bowland T, Ionescu P, Laster MJ, Weiskopf RB. The effect of anesthetic duration on kinetic and recovery characteristics of desflurane versus sevoflurane, and on the kinetic characteristics of compound A, in volunteers, 414
Ionescu P, Gong D. Circuit absorption of halothane, isoflurane, and sevoflurane, 1070
Koblin DD. A step toward consensus on general anesthesia (letter, reply), 446
see Gong DH
see Sonner J
see Steffey EP
Eisenach JC, see Owen MD
Elam M, see Ederberg S
Eledjam J-J, see Aya AGM
Ellis DS, see Glass JD
Ellison N. Anesthesia Education at Emory 1958–1985 (book review), 452
Emerson N, see Steffey EP
Enders-Klein C, see Dsida RM
Erd G, see Adelsmayr E
Eyraud D, Mouren S, Teugels K, Bertrand MA, Coriat P. Treating anesthesia-induced hypotension by angiotensin II in patients chronically treated with angiotensin-converting enzyme inhibitors, 259

Fajardo C, see Lema G
Faraday N, see Rosenfeld BA
Farnsworth ST, Gauthier ME, McJames SW, Zhang J, Mamalis N, Kopp CH. Ocular transmucosal absorption and toxicity of sufentanil in dogs, 138
Feeley TW. Perioperative Care: Anesthesia, Medicine and Surgery (book review), 922
Feletti F, see Imberti R
Felfernig M
see Blaicher AM
see Kettner SC
Ferreira JF, see Fleyfel M
Ferson D, see Brimacombe JR
Fields HL, see Robbins WR
Fife TJ, see Driver RP Jr
Filshie J, see McLure HA
Firestone L, see Banoub MF
Fish K, see Devitt JH
Fish P, see Devitt JH
Fisher HAG, see Groudine SB
Fishman RL, see Overdyk FJ
Fleyfel M, Provost N, Ferreira JF, Porte H, Bourzoufi K. Management of diaphragmatic hernia during pregnancy, 501
Forster A, see Klopfenstein CE
Förster H, see Behne M
Fortney JT, Gan TJ, Graczyk S, Wetchler B, Melson T, Khalil S, McKenzie R, Parrillo S, Glass PSA, Moote C,

Wermeling D, Parasuraman TV, Duncan B, Creed MR, S3A-409 and S3A-410 Study Groups. A comparison of the efficacy, safety, and patient satisfaction of ondansetron versus droperidol as antiemetics for elective outpatient surgical procedures, 731
Foss ML, see Owen MD
Foussat C, see Gueugniaud P-Y
François G, see Bruder N
Frank DU, see Horstman DJ
Franklin PD, see Beers RA
Frenette L, Cox J, McArdle P, Eckhoff D, Bynon S. Conjugated estrogen reduces transfusion and coagulation factor requirements in orthotopic liver transplantation, 1183
Fridrich P, see Sitzwohl C
Friedberg BL. Dissociative monitored anesthesia care not cited (letter), 1336
Frizelle HP, Moriarty DC. The "midhumeral" approach to the brachial plexus (letter), 447
Fu ES, see Naidu KA
Fu W, Stool LA, White PF, Husain MM. Is oral clonidine effective in modifying the acute hemodynamic response during electroconvulsive therapy?, 1127
Fujii Y, Toyooka H, Tanaka H. Prevention of postoperative nausea and vomiting with a combination of granisetron and droperidol, 613
Fujimori M, see Yabe M
Fujino Y, Nishimura M, Nishimura S, Taenaka N, Yoshiya I. Prolonged administration of isoflurane to patients with severe renal dysfunction, 440
Furuya H, see Kawaguchi M

Gajraj NM, see Chen YCJ
Gallagher JD. The effects of halothane on abnormal automaticity in canine cardiac Purkinje fibers, 488
Gallagher PR, see Cook-Sather SD
Galley HF, Dubbels AM, Webster NR. The effect of midazolam and propofol on interleukin-8 from human polymorphonuclear leukocytes, 1289
see Cuthbertson BH
Gan TJ, see Fortney JT
Gardner G, see Steinberg RB
Gäumann D, see Wilder-Smith OHG
Gauthier ME, see Farnsworth ST
Gellis JE, see Lynch EP
Gelman S. A step toward consensus on general anesthesia (letter), 446
Ischemic insult, kidney viability, and renal function (editorial), 1
Georgieff M
see Bothner U
see Schraag S
Gerancher JC, see Owen MD
Gerber H, see Konrad C
Gerlach E, see Dieterich H-JA
Gibson CS, see Reuben SS
Gielen MJM, see Visser WA
Gillin S, Sorkin LS. Gabapentin reverses the allodynia produced by the administration of anti-GD2 ganglioside, an immunotherapeutic drug, 111
Gilly H, see Michalek-Sauberer A
Gin T, Derrick JL, Chan MTV, Chui PT, Mak TWL. Postpartum patients have slightly prolonged neuromuscular block after mivacurium, 82
Glass JD, Ellis DS, Spiekermann BF, Hamill RJ. Intra- and postoperative airway management of a patient with a total laryngopharyngectomy and cervical esophagectomy (letter), 678
Glass PSA, see Fortney JT
Glosten B, see Mulroy M
Go G, see Sharrock NE
Godard J, see Gueugniaud P-Y
Goedecke R, see Latorre F
Goeters C, see Brodner G
Gogarten W, see Marcus MAE
Goldman L, see Lynch EP
Gombotz H, see Mahla E
Gomes JMA, see Lauretti GR
Gonano C, see Kettner SC
Gong D, see Eger EI II
Gong DH, Weiskopf RB, Neumann MA, Laster MJ, Eger EI II. In rats breathing from a nonrebreathing system, substitution of desflurane for isoflurane toward the end of anesthesia incompletely restores the time of recovery toward that of desflurane, 198
González AG, see Muñoz HAR
Goodman S, see Macario A
Goudsouzian NG, see Nishanian EV
Goyagi T, Tanaka M, Nishikawa T. Oral clonidine premedication reduces the awakening concentration of isoflurane, 410
Grabner CM, see Blaicher AM
Gracely R, see Sethna NF
Graczyk S, see Fortney JT
Graham MR, Thiessen DB, Mutch WAC. Left ventricular systolic and diastolic function is unaltered during propofol infusion in newborn swine, 717
Grant C
see Ludbrook GL
see Zheng D
Grass JA. Dose-response function of epidural fentanyl versus sufentanil (letter, reply), 919
Gratch DM. Critical Care Medicine in Transition: The Anesthesiologist's Role. Vol. 15, No. 4 of Anesthesiology Clinics of North America (book review), 1341
Gray AT, see Yost CS
Greenberg RS, see Kost-Byerly S
Greilich PE, see Whitten CW
Grenier B, Dubreuil M. Noninvasive monitoring of carbon dioxide: end-tidal versus transcutaneous carbon dioxide (letter), 675
Grimaudo V, see Alon E
Groen K, Mantel M, Zeijlmans PWM, Zeppenfeldt B, Olieman W, Stienstra R, van Kleef JW, Burm AGL. Pharmacokinetics of the enantiomers of bupivacaine and mepivacaine after epidural administration of the racemates, 361
Groudine SB, Fisher HAG, Kaufman RP Jr, Patel MK, Wilkins LJ, Mehta SA, Lumb PD. Intravenous lidocaine speeds the return of bowel function, decreases postoperative pain, and shortens hospital stay in patients undergoing radical retropubic prostatectomy, 235
Gründling M, see Hachenberg T
Gualea MR, see Imberti R
Gueugniaud P-Y, Abisseror M, Moussa M, Godard J, Foussat C, Petit P, Dodat H. The hemodynamic effects of pneumoperitoneum during laparoscopic surgery in healthy infants: assessment by continuous esophageal aortic blood flow echo-Doppler, 290
Gupta A, see Walldén J
Gupta R, Singhal SK, Rattan KN, Chhabra B. Management of congenital lobar emphysema with endobronchial intubation and controlled ventilation, 71
Guptill W, Carr D. Pain: Nociceptive and Neuropathic Mechanisms with Clinical Correlates. Vol. 15, No. 2 of Anesthesiology Clinics of North America (book review), 221
Gurevich B, see Talmor D

Hachenberg T, Karmann S, Pfeiffer B, Thomas H, Gründling M, Wendt M. The effect of dopexamine on ventilation perfusion distribution and pulmonary gas exchange in anesthetized, paralyzed patients, 314
Hague B, see Zhang J
Haisjackl M, Birnbaum JA, Redlin M, Schmutzler M, Waldenberger F, Lochs H, Konertz W, Kox W. Splanchnic oxygen transport and lactate metabolism during normothermic cardiopulmonary bypass in humans, 22
Hall DT, Varklet VL. Anesthesia: A Comprehensive Review (book review), 221
Hallén J, see Rawal N
Hamid SK, Selby IR, Sikich N, Lerman J. Vomiting after adenotonsillectomy in children: a comparison of ondansetron, dimenhydrinate, and placebo, 496
Hamill RJ, see Glass JD
Hammon JW Jr, see Butterworth JF IV
Hampson AJ, see Yost CS
Han Y-J
see Choe H
see Ko S-H
Hanaoka K, see Nishiyama T
Hara A, see Kokita N
Harrison P. Handbook of Palliative Care in Cancer (book review), 221
Harter RL, Kelly WB, Kramer MG, Perez CE, Dzwonczyk RR. A comparison of the volume and pH of gastric contents of obese and lean surgical patients, 147
Hartrick CT, Wise JJ, Patterson JS. Preemptive intrathecal ketamine delays mechanical hyperalgesia in the neuropathic rat, 557
Harvey SC, see Overdyk FJ
Hashizume H, see Kokita N
Hatano Y, see Yamamoto M
Haws JL, Herman N, Clark Y, Bjoraker R, Jones D. The chemical stability and sterility of sodium thiopental after preparation, 208
Heck M, see Boldt J
Hedman C, see Tarnow P
Hedman U, see Plesan A
Heidegger T, Kreienbühl G. Unsuccessful resuscitation under hypotensive epidural anesthesia during elective hip arthroplasty, 847
Heinen R, see Macario A
Heinrichs W
see Latorre F
see Weiler N
Helbo-Hansen HS, see Michelsen I
Henderson CL, Warriner CB, McEwen JA, Merrick PM. Intravenous regional anesthesia (letter, reply), 1334
Henthorn TK, see Dsida RM
Herbstman CH, Jaffee JB, Tuman KJ, Newman LM. An *in vivo* evaluation of four spinal needles used for the combined spinal-epidural technique, 520
Herman N, see Haws JL
Herrera E, see Alon E
Herrmann FAR, see Klopfenstein CE
Hettrick DA, see Pagel PS
Higgins MS, see DeLoach LJ
Hill GE, see Whitten CW
Hinder F, see Meyer JA
Hirsch M, see Schraag S
Ho ACY, see Tan PPC
Hoffmann VL, see Vercauteren MP
Högström H, see Curatolo M
Holthusen H, Stanton-Hicks M, Arndt JO. Sympathetic block does not reduce acute vascular pain in humans, 588
Hönemann CW, Brodner G, Van Aken H, Ruta U, Durieux ME, Möllhoff T. Aortic perforation during lumbar laminectomy, 493
Horie T, see Kubo N
Horlocker TT. Is continuous spinal anesthesia really so bad? (letter, reply), 215
The safety of continuous spinal anesthesia (letter, reply), 218
Wedel DJ. Spinal and epidural blockade and perioperative low molecular weight heparin: smooth sailing on the *Titanic* (editorial), 1153
Horne M, see Macario A
Horstman DJ, Frank DU, Rich GF. Prolonged inhaled NO attenuates hypoxic, but not monocrotaline-induced, pulmonary vascular remodeling in rats, 74
Horváth GA, see Klimscha W
Houltz E, see Ederberg S
House RM, see Katz JJ
Hrazdil J, see Lazar ER
Huang S, see Pang W-W
Huang YF, Pryor ME, Mather LE, Veering BT. Cardiovascular and central nervous system effects of intravenous levobupivacaine and bupivacaine in sheep, 797
see Mather LE
Hulstaert F, see Alon E
Husain MM, see Fu W
Hussong JW, see Eckmann DM
Hwang M-H, see Pang W-W
Hylander J, see Rawal N
Hynynen M, see Ranta SO-V

Ide T, Sakurai Y, Aono M, Nishino T. Minimum alveolar anesthetic concentrations for airway occlusion in cats: a new concept of minimum alveolar anesthetic concentration-airway occlusion response, 191
see Sakurai Y
Iida H
see Ishiyama T
see Ohata H
Ikeda K
see Kazama T
see Suzuki A
Ikuta Y, Shimoda O, Ushijima K, Terasaki H. Skin vasomotor reflex as an objective indicator to assess the level of regional anesthesia, 336
Illievich U, see Schramm WM
Illievich UM, see Sitzwohl C
Imberti R, Vairetti M, Gualea MR, Feletti F, Poma G, Richelmi P, Preseglio I, Bellomo G. The effects of thyroid hormone modulation on rat liver injury associated with ischemia-reperfusion and cold storage, 1187
Inoue T, see Shibata M
International Study of Perioperative Transfusion, see Bryson GL
Ionescu P
see Eger EI II
see Steffey EP
Iranami H, see Yamamoto M
Irarrazaval M, see Lema G
Irita K, Kawasaki T, Uenotsuchi T, Sakaguchi Y, Takahashi S. Does barbiturate therapy cause severe hypokalemia? (letter), 214
Ishimura N, see Kawaguchi M
Ishiyama T, Dohi S, Iida H. The vascular effects of topical and intravenous α_2-adrenoceptor agonist clonidine on canine pial microcirculation, 766
Ito Y
see Shakunaga K
see Yamazaki M
Ittleman FP, see Rathmell JP
Ivankovich AD, see McCarthy RJ
Ivy R, see Whitten CW
Ivy R III, see Patel SK

Jacka MJ, Sanderson F. Amyotrophic lateral sclerosis presenting during pregnancy, 542
Jaffee JB, see Herbstman CH
Jalil R, see Lema G

Jalkanen L, Rautoma P, Taivainen T, Meretoja OA. The pharmacodynamics of mivacurium preceded by atracurium or cisatracurium in children, 62
Jaquenod M, Rönnhedh C, Cousins MJ, Eckstein RP, Jordan V, Mather LE, Power I. Factors influencing ketorolac-associated perioperative renal dysfunction, 1090
Jaun N, see Boldt J
Joh J, Sila MK, Bastani B. Nondialyzability of fentanyl with high-efficiency and high-flux membranes (letter), 447
Johnson KJ, see Nader-Djalal N
Johnson MD, Schlanger M. Is miscarriage a risk factor for postdural puncture headache? (letter), 679
Jolly DT, see Lazar ER
Jomura K, see Shakunaga K
Jones D, see Haws JL
Jönsson A, see Tarnow P
Jordan V, see Jaquenod M
Joshi GP. Tourniquet release time and analgesic effects of intraarticular morphine (letter, reply), 1150
see Chen YCJ
see Song D
Joyner MJ, see Warner DO
Jun JH, Yaksh TL. The effect of intrathecal gabapentin and 3-isobutyl γ-aminobutyric acid on the hyperalgesia observed after thermal injury in the rat, 348

Kadoi Y. The differential effects of prostaglandin E_1 and nitroglycerin on regional cerebral oxygenation (letter, reply), 1339
Kagawa T, Maekawa N, Mikawa K, Nishina K, Yaku H, Obara H. The effect of halothane and sevoflurane on fatigue-induced changes in hamster diaphragmatic contractility, 392
Kähler CM, see Manfreda SE
Kahn RA, Weinberger J, Panah M. How selective is 7-nitroindazole, an inhibitor of neuronal nitric oxide synthase? (letter, reply), 679
Kalkman CJ, see Oei-Lim VLB
Kalso E, see Kontinen VK
Kanai Y, Katsuki H, Takasaki M. Graded, irreversible changes in crayfish giant axon as manifestations of lidocaine neurotoxicity *in vitro*, 569
Kanbak M. Tourniquet release time and analgesic effects of intraarticular morphine (letter), 1150
Kaneko S, see Morisaki H
Kapral S, see Marhofer P
Kapur PA, Steadman RH. Patient simulator competency testing: ready for takeoff? (editorial), 1157
Kardash KJ, see Wahba RWM
Karmann S, see Hachenberg T
Katchko L, see Talmor D
Kato H
see Kito K
see Mori K
Katoh T
see Kazama T
see Suzuki A
Katsuki H, see Kanai Y
Katz JJ, Mandell MS, House RM, Bilir BM, Barton B, Zamudio S. Cerebral blood flow velocity in patients with subclinical portal-systemic encephalopathy, 1005
Kaufman RP Jr, see Groudine SB
Kaufmann M, see Drop L
Kaushik S, see Ambesh SP
Kawaguchi M, Ishimura N, Kurehara K, Sakamoto T, Kitaguchi K, Furuya H. Mild hypothermia can attenuate nitroglycerin-induced vasodilation of pial arterioles in the cat, 546
Kawamata M, see Kawamata T
Kawamata T, Omote K, Kawamata M, Namiki A. Premedication with oral dextromethorphan reduces postoperative pain after tonsillectomy, 594
Kawasaki T, see Irita K
Kazama T, Ikeda K, Morita K, Katoh T, Kikura M. Propofol concentration required for endotracheal intubation with a laryngoscope or fiberscope and its interaction with fentanyl, 872
Keller C, Brimacombe J. Bronchial mucus transport velocity in paralyzed anesthetized patients: a comparison of the laryngeal mask airway and cuffed tracheal tube, 1280
see Adelsmayr E
Kellokumpu I, see Koivusalo A-M
Kelly DJ, Wallace DE, Brull SJ. Eyelid movement during complete neuromuscular block, 1332
Kelly PJ, see Patel PM
Kelly WB, see Harter RL
Kennedy TP, see Nader-Djalal N
Kersten JR, see Pagel PS
Kettner SC, Gonano C, Seebach F, Sitzwohl C, Acimovic S, Stark J, Schellongowski A, Blaicher A, Felfernig M, Zimpfer M. Endogenous heparin-like substances significantly impair coagulation in patients undergoing orthotopic liver transplantation, 691
see Blaicher AM
see Sitzwohl C
Khalil S, see Fortney JT
Khoury A, see Bouaziz H
Kikura M, see Kazama T
Kilaru PA, see Reuben SS
Kim D-C
see Choe H
see Ko S-H
Kim J-S, see Choe H
Kirsch JR, see Palmon SC
Kissin I, see Vinik HR
Kitaguchi K, see Kawaguchi M
Kito K, Kato H, Shibata M, Adachi T, Nakao S, Mori K. The effect of varied doses of epinephrine on duration of lidocaine spinal anesthesia in the thoracic and lumbosacral dermatomes, 1018
see Mori K
Klein W, see Mahla E
Klimscha W, Chiari A, Michalek-Sauberer A, Wildling E, Lerche A, Lorber C, Brinkmann H, Semsroth M. The efficacy and safety of a clonidine/bupivacaine combination in caudal blockade for pediatric hernia repair, 54
Horváth GA, Szikszay M, Dobos IA, Benedek GA. Antinociceptive effect of the S(+)-enantiomer of ketamine on carrageenan hyperalgesia after intrathecal administration in rats, 561
see Sitzwohl C
Klopfenstein CE, Herrmann FAR, Michel JP, Clergue FA, Forster A. The influence of an aging surgical population on the anesthesia workload: a ten-year survey, 1165
Knab JH, see Moore JM
Knight PR, see Nader-Djalal N
Ko S-H, Kim D-C, Han Y-J, Song H-S. Small-dose fentanyl: optimal time of injection for blunting the circulatory responses to tracheal intubation, 658
see Choe H
Koay CK, see Chow MYH
Koblin DD
see Eger EI II
see Yost CS
Koehntop DE, Beebe DS, Belani KG. Prophylactic use of dantrolene in a patient with central core disease (letter, reply), 915
Køhler F, see Michelsen I

Koidl C, see Mächler HE
Koivusalo A-M, Kellokumpu I, Scheinin M, Tikkanen I, Mäkisalo H, Lindgren L. A comparison of gasless mechanical and conventional carbon dioxide pneumoperitoneum methods for laparoscopic cholecystectomy, 153
Koizumi M, Matsumoto N, Ueda K. Influences of cardiopulmonary bypass and fentanyl anesthesia on hepatic circulation and oxygen metabolism in beagles, 1177
Kojima S, see Shakunaga K
Kokita N, Hara A, Abiko Y, Arakawa J, Hashizume H, Namiki A. Propofol improves functional and metabolic recovery in ischemic reperfused isolated rat hearts, 252
Konertz W, see Haisjackl M
Konrad C, Schüpfer G, Wietlisbach M, Gerber H. Learning manual skills in anesthesiology: is there a recommended number of cases for anesthetic procedures?, 635
Kontinen VK, Paananen S, Kalso E. The effects of the α_2-adrenergic agonist, dexmedetomidine, in the spinal nerve ligation model of neuropathic pain in rats, 355
Kopacz DJ, see Neal JM
Kopp CH, see Farnsworth ST
Kost-Byerly S, Tobin JR, Greenberg RS, Billett C, Zahurak M, Yaster M. Bacterial colonization and infection rate of continuous epidural catheters in children, 712
Kottra JA, see Litman RS
Kox W, see Haisjackl M
Kraft D, see Dieterich H-JA
Kramer DJ, see Neal JM
Kramer MG, see Harter RL
Kreienbühl G, see Heidegger T
Krishna YM, Reddy AR, Rao SM. Inspiratory collapse of alae nasi causing variable extrathoracic airway obstruction, 294
Kroin JS, see McCarthy RJ
Kroll DA. Anesthesia and Perioperative Care of the Combat Casualty (book review), 222
Kubo N, Sakuragi T, Mayama T, Shinokuma T, Horie T, Dan K. Collapsed anterior chamber during hypothermic cardiopulmonary bypass (letter), 919
Kühbacher C, see Mächler HE
Kulier A, see Trantina AE
Kumle B, see Boldt J
Kurehara K, see Kawaguchi M
Kurrek MM, see Devitt JH
Kussman BD, see O'Connor BR
Kwon J-Y, Bacher A, Zornow MH. Riluzole does not attenuate increases in hippocampal glutamate concentrations in a rabbit model of repeated transient global cerebral ischemia, 128

Labow RS, see Rosaeg OP
Laing DI, see Sher MH
Lalourcey L, see Aya AGM
Lam FK, see Lui P-W
Lambert DH. Is continuous spinal anesthesia really so bad? (letter), 214
Lang SA. Intravenous regional anesthesia (letter), 1334
  Ultrasound and the femoral three-in-one nerve block: weak methodology and inappropriate conclusions (letter), 1147
Laster MJ
  see Eger EI II
  see Gong DH
  see Steffey EP
Laterre P-FA, see De Kock M
Latorre F, Eberle B, Weiler N, Mienert R, Stanek A, Goedecke R, Heinrichs W. Laryngeal mask airway position and the risk of gastric insufflation, 867
  see Weiler N
Lau J, see Ballantyne JC
Laubenthal H, see Dieterich H-JA
Lauber R, see Curatolo M
Laupacis A, see Bryson GL
Lauretti GR, Reis MP, Mattos AL, Gomes JMA, Oliveira A-PM, Pereira NL. Epidural nonsteroidal antiinflammatory drugs for cancer pain, 117
Laurila R, see Ranta SO-V
Lazar ER, Jolly DT, Tam YK, Hrazdil J, Tawfik SR, Clanachan AS. Propofol and thiopental in a 1:1 volume mixture is chemically stable, 422
Lazor MA, see Lynch EP
Lee TH, see Swenson JD
Lee VC, Moscicki JC, DiFazio CA. Propofol sedation produces dose-dependent suppression of lidocaine-induced seizures in rats, 652
Legault C, see Butterworth JF IV
Leguerrier A, see Sicsic J-C
Lehzen S, see Palm S
Leimenstoll G, see Palm S
Lema G, Urzua J, Jalil R, Canessa R, Moran S, Sacco C, Medel J, Irarrazaval M, Zalaquett R, Fajardo C, Meneses G. Renal protection in patients undergoing cardiopulmonary bypass with preoperative abnormal renal function, 3
Lentschener C, Leveque JP, Mazoit JX, Benhamou D. The effect of pneumoperitoneum on intraocular pressure in rabbits with α-chymotrypsin–induced glaucoma, 1283
Leonoudakis D, see Yost CS
Lerche A, see Klimscha W
Lerman J, see Hamid SK
Lerut J, see De Kock M
Leveque JP, see Lentschener C
Levin A, Segal S, Datta S. Does combined spinal-epidural analgesia alter the incidence of paresthesia during epidural catheter placement? (letter), 448
Levine J, see Robbins WR
Levy WJ. Neuroprotection: A Society of Cardiovascular Anesthesiologists Monograph (book review), 921
Li C-Y, see Chou T-C
Li J, see Sonner J
Li L, see Xue FS
Liao X, see Xue FS
Lidegran G, see Rawal N
Liem TH, see Visser WA
Lin PJ, see Tan PPC
Lindahl SGE, see Sjögren D
Lindgren L, see Koivusalo A-M
Lipton JM, see Whitten CW
Lisbon A. Principles and Practice of Anesthesiology. 2nd Ed. (book review), 1340
  Nayyar P. The trauma airway: whose responsibility? (letter, reply), 1147
Lischke V, see Behne M
List WF
  see Mahla E
  see Manfreda SE
Litman RS, Kottra JA, Verga KA, Berkowitz RJ, Ward DS. Chloral hydrate sedation: the additive sedative and respiratory depressant effects of nitrous oxide, 724
Liu JH, see Xue FS
Liu K, see Chia Y-Y
Liu M, see Sethna NF
Liu SS
  see Moore JM
  see Neal JM
Liu X, see Chi OZ
Liu Y-C, see Chia Y-Y
Lloyd AT, see Palmon SC

Lobato EB, Paige GB, Brown MM, Bennett B, Davis JD. Pneumoperitoneum as a risk factor for endobronchial intubation during laparoscopic gynecologic surgery, 301
Lochs H, see Haisjackl M
Lockwood GG, see Ma D
Logeais Y, see Sicsic J-C
Lorber C, see Klimscha W
Lorenzen A-G, see Michelsen I
Lowe D, see Pagel PS
Lowenstein E, see Park KW
Lowes R, Mayhew JF. A suspected malignant hyperthermia episode during desflurane anesthesia (letter), 449
Ludbrook GL, Upton RN, Grant C, Martinez A. The effect of rate of administration on brain concentrations of propofol in sheep, 1301
see Zheng D
Lueger A, see Mächler HE
Luginbühl M, see Curatolo M
Lui P-W, Chan BCB, Chan FHY, Poon PWF, Wang H, Lam FK. Wavelet analysis of embolic heart sound detected by precordial Doppler ultrasound during continuous venous air embolism in dogs, 325
Lumb PD, see Groudine SB
Luo LK, see Xue FS
Lynch EP, Lazor MA, Gellis JE, Orav J, Goldman L, Marcantonio ER. The impact of postoperative pain on the development of postoperative delirium, 781
Lynn A, Nespeca MK, Bratton SL, Strauss SG, Shen DD. Clearance of morphine in postoperative infants during intravenous infusion: the influence of age and surgery, 958
Lynn AM, see Ramamoorthy C

Ma D, Wang C, Soo CKP, Chakrabarti MK, Lockwood GG, Whitwam JG. The effect of sevoflurane on spontaneous sympathetic activity, aδ and c somatosympathetic reflexes, and associated hemodynamic changes in dogs, 1079
Macaluso A, see Patel SK
Macario A, Horne M, Goodman S, Vitez T, Dexter F, Heinen R, Brown B. The effect of a perioperative clinical pathway for knee replacement surgery on hospital costs, 978
Mächler HE, Lueger A, Rehak P, Berger J, Veith W, Kühbacher C, Koidl C, Stark G, Metzler H. A new high-resolution esophageal electrocardiography recording technique: an experimental approach for the detection of myocardial ischemia, 34
see Trantina AE
Mackenzie C. Direct Laryngoscopy Video System (multimedia review), 921
Maddalozzo J, see Dsida RM
Maeda K, see Tsujimoto S
Maekawa N
see Kagawa T
see Nishina K
Mahla E, Rotman B, Rehak P, Atlee JL, Gombotz H, Berger J, List WF, Klein W, Metzler H. Perioperative ventricular dysrhythmias in patients with structural heart disease undergoing noncardiac surgery, 16
Mahlstedt D, see Bischoff P
Maier C, see Palm S
Maislin G, see Swan BA
Mak TWL, see Gin T
Mäkisalo H, see Koivusalo A-M
Makkes PC, see Oei-Lim VLB
Maland L, see Zhang J
Malviya S, see Tait AR
Mamalis N, see Farnsworth ST
Mandell MS, see Katz JJ
Manecke GR Jr, Poppers PJ. Esophageal stethoscope placement depth: its effect on heart and lung sound monitoring during general anesthesia, 1276
Manfreda SE, Dunzendorfer S, Schratzberger P, Buratti T, Reinisch N, Kähler CM, List WF, Wiedermann CJ. The chemotaxis of human peripheral blood B lymphocytes by β-endorphin is reversible by naloxone, 670
Mangin R, see Aya AGM
Mantel M, see Groen K
Marcantonio ER, see Lynch EP
Marcus MAE, Vertommen JD, Van Aken H, Gogarten W, Buerkle H. The effects of adding isoproterenol to 0.125% bupivacaine on the quality and duration of epidural analgesia in laboring parturients, 749
see Buerkle H
Marhofer P, Kapral S. Ultrasound and the femoral three-in-one nerve block: weak methodology and inappropriate conclusions (letter, reply), 1147
Markey JR, Montiague R, Winnie AP. The changing role of monitored anesthesia care in the ambulatory setting (letter, reply), 1335
Martin LD, see Tobin JR
Martinez A, see Ludbrook GL
Martinez AM, see Zheng D
Maruyama K, see Tsujimoto S
Mascia L, see McKeating EG
Mashimo T, see Abe K
Massa FC, see Campos JH
Masuda A, see Yamazaki M
Masuda J, see Morisaki H
Mather LE, Huang YF, Veering B, Pryor ME. Systemic and regional pharmacokinetics of levobupivacaine and bupivacaine enantiomers in sheep, 805
see Huang YF
see Jaquenod M
see Sharrock NE
Mathes DD. Treatment of myxedema coma for emergency surgery (letter), 450
Matsukawa T, see Nishiyama T
Matsumoto N, see Koizumi-M
Matsushima M, see Morisaki H
Mattos AL, see Lauretti GR
Maurer WG, see Whalley DG
Max MB, see Sethna NF
Maxson PM, see Warner ME
Mayama T, see Kubo N
Mayhew JF
see Abouleish AE
see Lowes R
see Medina KA
Mazoit JX, see Lentschener C
McArdle P, see Frenette L
McCarthy RJ, Kroin JS, Tuman KJ, Penn RD, Ivankovich AD. Antinociceptive potentiation and attenuation of tolerance by intrathecal co-infusion of magnesium sulfate and morphine in rats, 830
McEwen JA, see Henderson CL
McJames S,
see Swenson JD
see Farnsworth ST
McKeating EG, Andrews PJD, Mascia L. The relationship of soluble adhesion molecule concentrations in systemic and jugular venous serum to injury severity and outcome after traumatic brain injury, 759
McKenzie R, see Fortney JT
McLure HA, Filshie J. Radiological screening for Hickman catheter insertion (letter), 216
McShane AJ, see Doyle MT
Medel J, see Lema G
Medina KA, Mayhew JF. Generalized muscle rigidity and hypercarbia

with halothane and isoflurane, 297
Mehta SA, see Groudine SB
Melson T, see Fortney JT
Meneses G, see Lema G
Menestret P, see Sicsic J-C
Mercier FAAJ, Bouaziz HA, Benhamou D. Intrathecal analgesia for labor: how to take over with the epidural? (letter), 219
Mercier FJ, see Bouaziz H
Meretoja OA, see Jalkanen L
Merkind V, see Talmor D
Merrick PM, see Henderson CL
Mertes N, see Brodner G
Messana LD, see Ebert TJ
Metzler H
see Mächler HE
see Mahla E
Meyer JA, Theilmeier G, Van Aken H, Bone HG, Busse H, Waurick RA, Hinder F, Booke M. Inhaled prostaglandin E_1 for treatment of acute lung injury in severe multiple organ failure, 753
see Booke M
Meznik C, see Schmied H
Michalek-Sauberer A, Gilly H. A case of suspected malignant hyperthermia during desflurane administration (letter, reply), 916
Gilly H. A suspected malignant hyperthermia episode during desflurane anesthesia (letter, reply), 449
Gilly H. Prophylactic use of dantrolene in a patient with central core disease (letter), 915
see Klimscha W
see Schramm WM
Michel JP, see Klopfenstein CE
Michelsen I, Helbo-Hansen HS, Køhler F, Lorenzen A-G, Rydlund E, Bentzon MW. Prophylactic ephedrine attenuates the hemodynamic response to propofol in elderly female patients, 477
Michelsen LG. A more simple holder for the transesophageal echocardiography probe (letter), 677
Mienert R, see Latorre F
Mignat C, see Palm S
Mikawa K
see Kagawa T
see Nishina K
Mirza H, see Steinberg ES
Mizutani T, see Takahashi S
Mohl U, see Schraag S
Mok MS, see Pang W-W
Möllhoff T, see Hönemann CW
Molliex S, Montravers P, Dureuil B, Desmonts J-M. Halogenated anesthetics inhibit *Pseudomonas aeruginosa* growth in culture conditions reproducing the alveolar environment, 1075
Montiague R, see Markey JR
Montravers P, see Molliex S
Moore JM, Liu SS, Neal JM. Premedication with fentanyl and midazolam decreases the reliability of intravenous lidocaine test dose, 1015
Liu SS, Pollock JE, Neal JM, Knab JH. The effect of epinephrine on small-dose hyperbaric bupivacaine spinal anesthesia: clinical implications for ambulatory surgery, 973
see Neal JM
Moote C, see Fortney JT
Moran S, see Lema G
Mori K, Kato H, Stevens J, Kito K. Effects of epidural epinephrine on the spinal cord accumulation of epidural clonidine in rabbits (letter), 1145
see Kito K
Moriarty DC, see Frizelle HP
Morisaki H, Masuda J, Kaneko S, Matsushima M, Takeda J. Transient neurologic syndrome in one thousand forty-five patients after 3% lidocaine spinal anesthesia, 1023
Morita K, see Kazama T
Mort TC. Unplanned tracheal extubation outside the operating room: a quality improvement audit of hemodynamic and tracheal airway complications associated with emergency tracheal reintubation, 1171
Moscicki JC, see Lee VC
Mosteller F, see Ballantyne JC
Mouren S, see Eyraud D
Moussa M, see Gueugniaud P-Y
Mugawar M, Rajender Y, Purohit AK, Sastry RA, Sundaram C, Rammurti S. Anesthetic management of Von Hippel-Lindau syndrome for excision of cerebellar hemangioblastoma and pheochromocytoma surgery, 673
Muhm M, Sunder-Plassmann G, Apsner R. Radiological screening for Hickman catheter insertion (letter, reply), 216
Müller H, see Trantina AE
Mullin V, de Rosayro M, Quint D. Mechanism of action caudal steroids for piriformis syndrome (letter), 680
Mulroy M, Glosten B. The epinephrine test dose in obstetrics: note the limitations (editorial), 923
Mulroy MF. Regional anesthesia techniques for outpatient shoulder arthroscopy (letter), 217
Mund K, see Boldt J
Munn SR, see Warner ME
Muñoz HAR, González AG, Dagnino JA. Cardiac output and onset time of different muscle relaxants (letter, reply), 1146
Munro HM, see Tait AR
Muravchick S. Management of the Elderly Surgical Patient. Vol. 9, No. 4 of Problems in Anesthesia (book review), 1152
Murphy FL. The changing role of monitored anesthesia care in the ambulatory setting (letter), 1335
Mutch WAC, see Graham MR
Myles PS. Case-control and cohort studies in anesthesia (letter), 1149
Mythen MG, see Bennett-Guerrero E

Nader-Djalal N, Knight PR, Bacon MF, Tait AR, Kennedy TP, Johnson KJ. Alterations in the course of acid-induced lung injury in rats after general anesthesia: volatile anesthetics versus ketamine, 141
Nagaraj L, see Colonna-Romano P
Naidu KA, Fu ES, Prockop LD. Effects of epidural epinephrine on the spinal cord accumulation of epidural clonidine in rabbits (letter, reply), 1145
Nakao S. Anesthesia for Cardiac Surgery (book review), 1340
see Kito K
Namiki A
see Kawamata T
see Kokita N
Narchi P, see Bouaziz H
Nayyar P, see Lisbon A
Neal JM, Moore JM, Kopacz DJ, Liu SS, Kramer DJ, Plorde JJ. Quantitative analysis of respiratory, motor, and sensory function after supraclavicular block, 1239
see Moore JM
Nespeca MK, see Lynn A
Neumann MA, see Gong DH
Newman LM, see Herbstman CH
Newman MF, Croughwell ND, White WD, Sanderson I, Spillane W, Reves JG. Pharmacologic electroencephalographic suppression during

cardiopulmonary bypass: a comparison of thiopental and isoflurane, 246
Nguyen N-D, see Rosenfeld BA
Nicolson SC, see Cook-Sather SD
Nikonorov IM, Blanck TJJ, Recio-Pinto E. The effects of halothane on single human neuronal L-type calcium channels, 885
Nishanian EV, Goudsouzian NG. Carbon dioxide embolism during hip arthography in an infant, 299
Nishikawa K, see Yabe M
Nishikawa T
see Goyagi T
see Tanaka M
Nishimura M, see Fujino Y
Nishimura S, see Fujino Y
Nishina K, Akamatsu H, Mikawa K, Shiga M, Maekawa N, Obara H, Niwa Y. The inhibitory effects of thiopental, midazolam, and ketamine on human neutrophil functions, 159
see Kagawa T
Nishino T
see Ide T
see Sakurai Y
Nishiyama T, Matsukawa T, Hanaoka K. The effects of age and gender on the optimal premedication dose of intramuscular midazolam, 1103
Niu S, see Zhang J
Niwa Y, see Nishina K
Noel AG, see Devitt JH
Norman PH, Coveler LA, Daley MD, Dugas MJ. Failure of a laryngoscope blade (letter), 448
North FP, see Driver RP Jr
Nottberg H, see Brodner G
Nozaki-Taguchi N, Yamamoto T. The interaction of FK409, a novel nitric oxide releaser, and peripherally administered morphine during experimental inflammation, 367

Obara H
see Kagawa T
see Nishina K
Oberwalder P, see Trantina AE
Oei-Lim VLB, Kalkman CJ, Makkes PC, Ooms WG. Patient-controlled versus anesthesiologist-controlled conscious sedation with propofol for dental treatment in anxious patients, 967
Ogawa K, see Yamamoto M
Oh Y-S, see Bahk J-H
Ohata H, Iida H, Watanabe Y, Dohi S. The optimal test dose of epinephrine for epidural injection with lidocaine solution in awake patients premedicated with oral clonidine, 1010
Olieman W, see Groen K
Oliveira A-PM, see Lauretti GR
Omote K, see Kawamata T
Onaran O, see Adsan H
Ooms WG, see Oei-Lim VLB
Orav J, see Lynch EP
Orser B, Oxorn D. The effects of midazolam in propofol-induced anesthesia (letter, reply), 1148
Overand PT, Teply JF. Vasopressin for the treatment of refractory hypotension after cardiopulmonary bypass, 1207
Overdyk FJ, Harvey SC, Fishman RL, Shippey F. Successful strategies for improving operating room efficiency at academic institutions, 896
Owen MD, D'Angelo R, Gerancher JC, Thompson JM, Foss ML, Babb JD, Eisenach JC. 0.125% ropivacaine is similar to 0.125% bupivacaine for labor analgesia using patient-controlled epidural infusion, 527
Owens TM, see Doyle MT
Oxorn D, see Orser B
O'Connor BR, Kussman BD, Park KW. Severe hypercarbia during cardiopulmonary bypass: a complication of CO_2 flooding of the surgical field, 264
O'Hara P, see Sprung J
O'Leary CE, see Beers RA

Paananen S, see Kontinen VK
Pagel PS, Hettrick DA, Kersten JR, Tessmer JP, Lowe D, Warltier DC. Etomidate adversely alters determinants of left ventricular afterload in dogs with dilated cardiomyopathy, 932
Paige GB, see Lobato EB
Palm S, Lehzen S, Mignat C, Steinmann J, Leimenstoll G, Maier C. Does prolonged oral treatment with sustained-release morphine tablets influence immune function?, 166
Palmon SC, Lloyd AT, Kirsch JR. The effect of needle gauge and lidocaine pH on pain during intradermal injection, 379
Panah M, see Kahn RA
Pandit UA, see Tait AR
Pang W-W, Mok MS, Huang S, Hwang M-H. The analgesic effect of fentanyl, morphine, meperidine, and lidocaine in the peripheral veins: a comparative study, 382
Papadakos PJ. Handbook of Mechanical Ventilatory Support, 2nd Ed. (book review), 222
Papousek A, see Schramm WM
Pappagallo M, see Robbins WR
Parasuraman TV, see Fortney JT
Park KW, Dai HB, Lowenstein E, Seilke FW. Epithelial dependence of the bronchodilatory effect of sevoflurane and desflurane in rat distal bronchi, 646
see O'Connor BR
Parrillo S, see Fortney JT
Patel MK, see Groudine SB
Patel PM, Drummond JC, Cole DJ, Kelly PJ, Watson M. Isoflurane and pentobarbital reduce the frequency of transient ischemic depolarizations during focal ischemia in rats, 773
Patel SK, Whitten CW, Ivy R III, Macaluso A, Pennant JH. Failure of the laryngeal mask airway: an undiagnosed laryngeal carcinoma, 438
Patterson JS, see Hartrick CT
Pellissier D, see Bruder N
Penn RD, see McCarthy RJ
Pennant JH. The Laryngeal Mask Airway: A Review and Practical Guide (book review), 452
see Patel SK
Peray P, see Capdevila X
Pereira NL, see Lauretti GR
Perez CE, see Harter RL
Perez-Vela JL,
see Sanchez-Izquierdo-Riera JA
Peter K, see Dieterich H-JA
Petersen-Felix S, see Curatolo M
Petit P, see Gueugniaud P-Y
Petit PL, see Branche PE
Pfeiffer B, see Hachenberg T
Pham-Dang C. Sternocleidomastoid muscle as surface landmark for supraclavicular nerve blocks (letter, reply), 1150
Philippi W, see Striebel HW
Pivalizza EG, see Campos CJ
Plesan A, Hedman U, Xu X-J, Wiesenfeld-Hallin Z. Comparison of ketamine and dextromethorphan in potentiating the antinociceptive effect of morphine in rats, 825
Plorde JJ, see Neal JM
Pogatzki E, see Brodner G
Poirier T, see Bouaziz H
Polaner DM. Airway Management in Pediatric Anesthesia. Vol. 35, No. 3 of International Anesthesiology Clinics (book review), 921

Pollock JE, see Moore JM
Pölz W, see Dieterich H-JA
Poma G, see Imberti R
Poon PWF, see Lui P-W
Poppers PJ, see Manecke GR Jr
Porte H, see Fleyfel M
Pothula SNM, see Bhargava M
Power I, see Jaquenod M
Prado AA, see Cardoso MAMSC
Preseglio I, see Imberti R
Prielipp RC, see Rathmell JP
Prockop LD, see Naidu KA
Provost N, see Fleyfel M
Pryor ME
see Huang YF
see Mather LE
Purdy EP, Ajimal GS. Vision loss after lumbar epidural steroid injection, 119
Purohit AK, see Mugawar M

Qi J, see Tang J
Quint D, see Mullin V
Quintin L, see Bruandet N

Rady MY. Case-control and cohort studies in anesthesia (letter, reply), 1149
Rajender Y, see Mugawar M
Ramamoorthy C, Anderson GD, Williams GD, Lynn AM. Pharmacokinetics and side effects of milrinone in infants and children after open heart surgery, 283
Rammurti S, see Mugawar M
Ranta SO-V, Laurila R, Saario J, Ali-Melkkilä T, Hynynen M. Awareness with recall during general anesthesia: incidence and risk factors, 1084
Rao SM, see Krishna YM
Rathmell JP, Prielipp RC, Butterworth JF, Williams E, Villamaria F, Testa L, Viscomi C, Ittleman FP, Baisden CE, Royster RL. A multicenter, randomized, blind comparison of amrinone with milrinone after elective cardiac surgery, 683
Rattan KN, see Gupta R
Rautoma P, see Jalkanen L
Rawal N, Axelsson K, Hylander J, Allvin RA, Amilon CA, Lidegran G, Hallén J. Postoperative patient-controlled local anesthetic administration at home, 86
Recio-Pinto E, see Nikonorov IM
Reddy AR, see Krishna YM
Redlin M, see Haisjackl M
Rehak P
see Mächler HE
see Mahla E
Reichenthal E, see Talmor D
Reiner A, see Zagvazdin Y
Reinisch N, see Manfreda SE
Reinprecht A, see Sitzwohl C
Reis MP, see Lauretti GR
Reiter WJ, see Blaicher AM
Rentero N, see Bruandet N
Reuben SS, Steinberg RB, Cohen MA, Kilaru PA, Gibson CS. Intraarticular morphine in the multimodal analgesic management of postoperative pain after ambulatory anterior cruciate ligament repair, 374
see Steinberg RB
Reves JG, see Newman MF
Rich GF, see Horstman DJ
Richardson MG, Dooley JW. The effects of general versus epidural anesthesia for outpatient extracorporeal shock wave lithotripsy, 1214
Richelmi P, see Imberti R
Ricksten S-E, see Ederberg S
Rifat KR, see Wilder-Smith OHG
Rigler B, see Trantina AE
Riley RH, Wong JWF. Retrieval of air from the arrow antecubital catheter (letter), 1337
Rimbäck G, see Tarnow P
Ritchie E, see Tong D
Ritman EL, see Warner DO
Ritter L, see Alon E
Robbins WR, Staats PS, Levine J, Fields HL, Allen RW, Campbell JN, Pappagallo M. Treatment of intractable pain with topical large-dose capsaicin: preliminary report, 579
Robertson S, see Zhang J
Rönnhedh C, see Jaquenod M
Rosaeg OP, Cicutti NJ, Labow RS. The effect of oxytocin on the contractile force of human atrial trabeculae, 40
Rosenfeld BA, Nguyen N-D, Sung I, Faraday N. Neuroendocrine stress hormones do not recreate the postoperative hypercoagulable state, 640
Rosenow DE, Albrechtsen M, Stolke D. A comparison of patient-controlled analgesia with lornoxicam versus morphine in patients undergoing lumbar disk surgery, 1045
Rosenthal JA. Diaphragmatic paralysis complicating alcohol splanchnic nerve block, 845
Rotman B, see Mahla E
Royster RL
see Butterworth JF IV
see Rathmell JP
Rubenovitch J, see Capdevila X
Ruetsch YA, see Borgeat A
Ruta U, see Hönemann CW
Rydlund E, see Michelsen I

S3A-409 S3A-410 Study Groups, see Fortney JT
Saario J, see Ranta SO-V
Sacco C, see Lema G
Sagnard PE, see Branche PE
Sakaguchi Y, see Irita K
Sakamoto T, see Kawaguchi M
Sakuragi T, see Kubo N
Sakurai Y, Ide T, Aono M, Nishino T. The inhibitory influence of pulmonary vagal afferents on respiratory distress induced by airway occlusion in halothane-anesthetized cats, 398
see Ide T
Saldien V, see Vercauteren MP
Sanansilp V, Areewatana S, Tonsukchai N. Droperidol and the side effects of epidural morphine after cesarean section, 532
Sanchez-Izquierdo-Riera JA, Caballero-Cubedo RE, Perez-Vela JL, Ambros-Checa A, Cantalapiedra-Santiago JA, Alted-Lopez E. Propofol versus midazolam: safety and efficacy for sedating the severe trauma patient, 1219
Sanderson F, see Jacka MJ
Sanderson I, see Newman MF
Sá Rêgo Mônica M, White PF. Dissociative monitored anesthesia care not cited (letter, reply), 1336
see White PF
Sarna MC. Intravenous oxytocin in patients undergoing elective cesarean section (letter, reply), 1334
Sasaki R, see Yamazaki M
Sastry RA, see Mugawar M
Sata T, see Aoyama K
Sato S, see Takahashi S
Satone T, see Shakunaga K
Satumae T, see Watanabe S
Saunders TA, see Steinberg ES
Scaramozzino P, see Curatolo M
Scheinin M, see Koivusalo A-M
Scheitza W, see Striebel HW
Schellongowski A, see Kettner SC
Schiferer A, see Schmied H
Schimetta W, see Dieterich H-JA
Schlanger M, see Johnson MD
Schmied H, Schiferer A, Sessler DI, Meznik C. The effects of red-cell scavenging, hemodilution, and

active warming on allogenic blood requirements in patients undergoing hip or knee arthroplasty, 387
Schmutzler M, see Haisjackl M
Schoenwald PK, see Sprung J
Schraag S, Mohl U, Hirsch M, Stolberg E, Georgieff M. Recovery from opioid anesthesia: the clinical implication of context-sensitive half-times, 184
Schramm WM, Papousek A, Michalek-Sauberer A, Czech T, Illievich U. The cerebral and cardiovascular effects of cisatracurium and atracurium in neurosurgical patients, 123
Schratzberger P, see Manfreda SE
Schreiner MS, see Cook-Sather SD
Schroeder DR, see Warner ME
Schulte am Esch J, see Bischoff P
Schulzki C, see Brodner G
Schüpfer G, see Konrad C
Schwinn DA, see Bennett-Guerrero E
Sculco TP, see Sharrock NE
Seebach F, see Kettner SC
Segal S, Csavoy AN, Datta S. Placental tissue enhances uterine relaxation by nitroglycerin, 304
see Levin A
Selby IR, see Hamid SK
Seligman I, see Eckmann DM
Sellke FW, see Park KW
Semsroth M, see Klimscha W
Serre-Cousine O, see Capdevila X
Sessler DI, see Schmied H
Sethna NF, Liu M, Gracely R, Bennett GJ, Max MB. Analgesic and cognitive effects of intravenous ketamine-alfentanil combinations versus either drug alone after intradermal capsaicin in normal subjects, 1250
Shakunaga K, Kojima S, Jomura K, Shimizu Y, Satone T, Ito Y. Ketamine suppresses the production and release of endothelin 1 from cultured bovine endothelial cells, 1098
Shapira Y, see Talmor D
Sharrock NE, Mather LE, Go G, Sculco TP. Arterial and pulmonary arterial concentrations of the enantiomers of bupivacaine after epidural injection in elderly patients, 812
Shen DD, see Lynn A
Sher MH, Laing DI, Brands E. Life-threatening upper airway obstruction after glossopharyngeal nerve block: possibly due to an inappropriately large dose of bupivacaine? (letter), 678
Shibata M, Wakisaka S, Inoue T, Shimizu T, Yoshiya I. The effect of electroconvulsive treatment on thermal hyperalgesia and mechanical allodynia in a rat model of peripheral neuropathy, 584
see Kito K
Shiga M, see Nishina K
Shigematsu A, see Aoyama K
Shime N, Sugimoto E. Lumbar sympathetic ganglion block in a patient with painful legs and moving toes syndrome, 1056
Shimizu T, see Shibata M
Shimizu Y, see Shakunaga K
Shimoda O, see Ikuta Y
Shin Y-S, see Ahn EK
Shinokuma T, see Kubo N
Shippey F, see Overdyk FJ
Shorten GD, see Sieber TJ
Sia ATH, see Chow MYH
Sicsic J-C, Duranteau J, Corbineau HA, Antoun S, Menestret P, Sitbon P, Leguerrier A, Logeais Y, Ecoffey C. Gastric mucosal oxygen delivery decreases during cardiopulmonary bypass despite constant systemic oxygen delivery, 455
Sieber TJ, Zbinden AM, Curatolo M, Shorten GD. Tracheal intubation with rocuronium using the "timing principle", 1137
see Curatolo M
Siewert M, see Tait AR
Sikich N, see Hamid SK
Sila MK, see Joh J
Silbergleit R, Walls RM. The trauma airway: whose responsibility? (letter), 1146
Simhi E, Brandom BW. Cardiac output and onset time of different muscle relaxants (letter), 1146
Simon AE, see Berman MF
Singh H. Intrathecal fentanyl with small-dose dilute bupivacaine: better anesthesia without prolonging recovery (letter), 917
Singhal SK, see Gupta R
Sirtl C, see Dieterich H-JA
Sitbon P, see Sicsic J-C
Sitzwohl C, Kettner SC, Reinprecht A, Dietrich W, Klimscha W, Fridrich P, Sladen RN, Illievich UM. The arterial to end-tidal carbon dioxide gradient increases with uncorrected but not with temperature-corrected $Paco_2$ determination during mild to moderate hypothermia, 1131
see Kettner SC
Sjögren D, Lindahl SGE, Sollevi A. Ventilatory responses to acute and sustained hypoxia during isoflurane anesthesia, 403
Sladen RN, see Sitzwohl C
Slaughter TF, see Dalamangas LC
Snyder IS, see Driver RP Jr
Sollevi A, see Sjögren D
Song D, Joshi GP, White PF. Fast-track eligibility after ambulatory anesthesia: a comparison of desflurane, sevoflurane, and propofol, 267
Song H-S
see Choe H
see Ko S-H
Sonner J, Li J, Eger EI II. Desflurane and nitrous oxide, but not nonimmobilizers, affect nociceptive responses, 629
Soo CKP, see Ma D
Sorkin LS, see Gillin S
Souter MJ, Andrews PJD, Alston RP. Propofol does not ameliorate cerebral venous oxyhemoglobin desaturation during hypothermic cardiopulmonary bypass, 926
Spahn DR. The differential effects of prostaglandin E_1 and nitroglycerin on regional cerebral oxygenation (letter), 1338
Sperry RJ. Anesthesia for the Patient with Neurologic Disease. Anesthesiology Clinics of North America. Vol. 15, No. 3 (book review), 453
Spiekermann BF, see Glass JD
Spillane W, see Newman MF
Sprung J, Abdelmalak B, Schoenwald PK. Vasovagal cardiac arrest during the insertion of an epidural catheter and before the administration of epidural medication, 1263
Correia R, Schoenwald PK, Bhambani R, O'Hara P. A successful continuous spinal anesthetic in a patient with preoperative baseline hypercapnia exceeding 100 mm Hg, 591
see Banoub MF
Staacke T. Ethics of human toxicity study questioned (letter), 445
Staacke TS, see Ebert TJ
Staats PS, see Robbins WR
Standl T. The safety of continuous spinal anesthesia (letter), 218
Stanek A, see Latorre F
Stanley TH, see Zhang J
Stanton-Hicks M, see Holthusen H
Stark G, see Mächler HE
Stark J, see Kettner SC

Steadman RH, see Kapur PA
Steffey EP, Laster MJ, Ionescu P, Eger EI II, Emerson N. Ventilatory effects of the nonimmobilizer 1,2-dichlorohexafluorocyclobutane (2N) in swine, 173
Steinberg ES, Adsumelli RSN, Saunders TA, Mirza H. Successful epidural anesthesia for cesarean section for sextuplets, 1236
Steinberg RB, Reuben SS, Gardner G. The dose-response relationship of ketorolac as a component of intravenous regional anesthesia with lidocaine, 791
see Reuben SS
Steinbrook RA. Epidural anesthesia and gastrointestinal motility, 837
Steinmann J, see Palm S
Stevens J, see Mori K
Stevens JB, Wheatley LD. Tracheal intubation in ambulatory surgery patients: using remifentanil and propofol without muscle relaxants, 45
Stienstra R
see Groen K
see van den Nieuwenhuyzen MCO
Stiff JL, see DeLoach LJ
Stolberg E, see Schraag S
Stolke D, see Rosenow DE
Stool LA, see Fu W
Strauss SG, see Lynn A
Streisand JB, see Zhang J
Striebel HW, Scheitza W, Philippi W, Behrens U, Toussaint S. Quantifying oral analgesic consumption using a novel method and comparison with patient-controlled intravenous analgesic consumption, 1051
Suarez T, see Ballantyne JC
Sugimoto E, see Shime N
Sugiyama T, see Tsujimoto S
Sun HL. Naloxone-precipitated acute opioid withdrawal syndrome after epidural morphine, 544
Sundaram C, see Mugawar M
Sunder-Plassmann G, see Muhm M
Sung I, see Rosenfeld BA
Suzuki A, Katoh T, Ikeda K. The effect of adenosine triphosphate on sevoflurane requirements for minimum alveolar anesthetic concentration and minimum alveolar anesthetic concentration-awake, 179
Svensson S-E, see Ederberg S
Swan BA, Maislin G, Traber KB. Symptom distress and functional status changes during the first seven days after ambulatory surgery, 739
Swenson JD, Lee TH, McJames S. The effect of prior dural puncture on cerebrospinal fluid sufentanil concentrations in sheep after the administration of epidural sufentanil, 794
Swisher JL, see Bogetz MS
Szalai J-P, see Devitt JH
Szikszay M, see Klimscha W

Taenaka N, see Fujino Y
Taguchi N, see Watanabe S
Tait AR, Pandit UA, Voepel-Lewis T, Munro HM, Malviya S. Use of the laryngeal mask airway in children with upper respiratory tract infections: a comparison with endotracheal intubation, 706
Voepel-Lewis T, Siewert M, Malviya S. Factors that influence parents' decisions to consent to their child's participation in clinical anesthesia research, 50
see Nader-Djalal N
Taivainen T, see Jalkanen L
Tajima T, see Yamamoto M
Takada K, see Abe K
Takahashi S, Mizutani T, Sato S. Changes in oxygen consumption and carbon dioxide elimination after tourniquet release in patients breathing spontaneously under epidural anesthesia, 90
see Irita K
Takasaki M, see Kanai Y
Takeda J, see Morisaki H
Takenaka I, see Aoyama K
Takeshima R, see Watanabe S
Talmor D, Shapira Y, Artru AA, Gurevich B, Merkind V, Katchko L, Reichenthal E. 0.45% saline and 5% dextrose in water, but not 0.9% saline or 5% dextrose in 0.9% saline, worsen brain edema two hours after closed head trauma in rats, 1225
Tam YK, see Lazar ER
Tan PPC, Chu J-J, Ho ACY, Cheng K-S, Lin PJ, Chang C-H. A modified endotracheal tube for infants and small children undergoing video-assisted thoracoscopic surgery, 1212
Tanaka H, see Fujii Y
Tanaka M, Nishikawa T. Simulation of an epidural test dose with intravenous epinephrine in sevoflurane-anesthetized children, 952
Nishikawa T. The combination of epinephrine and isoproterenol as a simulated epidural test dose in isoflurane-anesthetized adults, 1312
see Goyagi T
Tang J, Wang B, White PF, Watcha MF, Qi J, Wender RH. The effect of timing of ondansetron administration on its efficacy, cost-effectiveness, and cost-benefit as a prophylactic antiemetic in the ambulatory setting, 274
Tanus dos Santos JAE. Propofol and the supraventricular tachydysrhythmias in children (letter), 914
Tarnow P, Cassuto J, Jönsson A, Rimbäck G, Hedman C. Postoperative analgesia by D-*myo*-inositol-1,2,6-trisphosphate in patients undergoing cholecystectomy, 107
Tassonyi EA, see Wilder-Smith OHG
Tawfik SR, see Lazar ER
Teply JF, see Overand PT
Terai T, see Yabe M
Terasaki H, see Ikuta Y
Tessler MJ, see Wahba RWM
Tessmer JP, see Pagel PS
Testa L, see Rathmell JP
Teugels K, see Eyraud D
Theilmeier G
see Booke M
see Meyer JA
Thiessen DB, see Graham MR
Thomas H
see Behne M
see Hachenberg T
Thompson JM, see Owen MD
Tikkanen I, see Koivusalo A-M
Tobin JR, Martin LD. Eleventh annual meeting of the Society for Pediatric Anesthesia, San Diego, California, October 17, 1997, 909
see Kost-Byerly S
Tong D, Chung F, Ritchie E. Regional anesthesia techniques for outpatient shoulder arthroscopy (letter, reply), 217
Tong SY, see Xue FS
Tonsukchai N, see Sanansilp V
Toussaint S, see Striebel HW
Toyooka H, see Fujii Y
Traber KB, see Swan BA
Trantina AE, Mächler H, Kulier A, Bergmann P, Dacar D, Müller H, Oberwalder P, Zenker G, Rigler B. Preoperative prostaglandin E_1 therapy in a patient with atrial septal defect and predominant right-to-left shunting, 703
Tsujimoto S, Maeda K, Sugiyama T, Yokochi A, Chikusa H, Maruyama K. Efficacy of prolonged large-dose

dantrolene for severe neuroleptic malignant syndrome, 1143
Tulloch HV, see Cook-Sather SD
Tulunay M, see Adsan H
Tuman KJ
see Herbstman CH
see McCarthy RJ
Twomey CF, see Doyle MT

Ueda K, see Koizumi M
Uenotsuchi T, see Irita K
Uhrich TD, see Ebert TJ
Upton RN
see Ludbrook GL
see Zheng D
Urzua J, see Lema G
Ushijima K, see Ikuta Y

Vairetti M, see Imberti R
Van Aken H
see Booke M
see Buerkle H
see Hönemann CW
see Marcus MAE
see Meyer JA
van den Nieuwenhuyzen MCO, Stienstra R, Burm AGL, Vletter AA, van Kleef JW. Alfentanil as an adjuvant to epidural bupivacaine in the management of postoperative pain after laparotomies: lack of evidence of spinal action, 574
van Egmond J, see Visser WA
van Kleef JW
see Groen K
see van den Nieuwenhuyzen MCO
Van Obbergh L, see De Kock M
Varklet VL, see Hall DT
Veering BT
see Huang YF
see Mather LE
Veith W, see Mächler HE
Vercauteren MP, Coppejans HC, Hoffmann VL, Saldien V, Adriaensen HA. Small-dose hyperbaric versus plain bupivacaine during spinal anesthesia for cesarean section, 989
Verga KA, see Litman RS
Verghese C, see Brimacombe JR
Vertommen JD, see Marcus MAE
Viale J-P, see Branche PE
Villamaria F, see Rathmell JP
Vinik HR. The effects of midazolam in propofol-induced anesthesia (letter), 1148
Kissin I. Rapid development of tolerance to analgesia during remifentanil infusion in humans, 1307
Viscomi C, see Rathmell JP
Viscomi CM. Intrathecal analgesia for labor: how to take over with the epidural? (letter, reply), 219
Visser WA, Liem TH, van Egmond J, Gielen MJM. Extension of sensory blockade after thoracic epidural administration of a test dose of lidocaine at three different levels, 332
Vitez T, see Macario A
Vizquel L, Barthe A, Benani MR. Sternocleidomastoid muscle as surface landmark for supraclavicular nerve blocks (letter), 1149
Vletter AA, see van den Nieuwenhuyzen MCO
Voepel-Lewis T, see Tait AR
Vogt NH, see Bothner U

Wahba RWM, Tessler MJ, Kardash KJ. Carbon dioxide tensions during anesthesia in the prone position, 668
Wakeling HG. Laryngeal mask insertion techniques (letter, reply), 1338
Wakisaka S, see Shibata M
Waldenberger F, see Haisjackl M
Walker JS, Carmody JJ. Experimental pain in healthy human subjects: gender differences in nociception and in response to ibuprofen, 1257
Wallace DE, see Kelly DJ
Walldén J, Gupta A, Carlsén H-O. Supraventricular tachycardia induced by Datex patient monitoring system (letter), 1339
Walls RM, see Silbergleit R
Wang B, see Tang J
Wang C, see Ma D
Wang H, see Lui P-W
Ward DS, see Litman RS
Warltier DC, see Pagel PS
Warner DO, Warner MA, Joyner MJ, Ritman EL. The effect of nitrous oxide on chest wall function in humans and dogs, 1058
Warner MA
see Warner DO
see Warner ME
Warner ME, Contreras MG, Warner MA, Schroeder DR, Munn SR, Maxson PM. Diabetes mellitus and difficult laryngoscopy in renal and pancreatic transplant patients, 516
Warriner CB, see Henderson CL
Watanabe S, Satumae T, Takeshima R, Taguchi N. Opisthotonos after flumazenil administered to antagonize midazolam previously administered to treat developing local anesthetic toxicity (letter), 677
Watanabe Y, see Ohata H
Watcha MF, see Tang J
Watson M, see Patel PM
Watts JC. Latex allergy and anesthesia (letter), 919
Waurick RA, see Meyer JA
Webster NR
see Cuthbertson BH
see Galley HF
Wedel DJ, see Horlocker TT
Weiler N, Latorre F, Eberle B, Heinrichs W. Gastric insufflation and the laryngeal mask (letter, reply), 914
see Latorre F
Weinberger J, see Kahn RA
Weiskopf RB
see Eger EI II
see Gong DH
Weiss HR, see Chi OZ
Wells GA, see Bryson GL
Wender RH, see Tang J
Wendt M, see Hachenberg T
Wermeling D, see Fortney JT
Westerlind A, see Ederberg S
Wetchler B, see Fortney JT
Whalley DG, Maurer WG. Hemodynamic effects of anesthesia in patients receiving angiotensin-converting enzyme inhibitors (letter), 215
Wheatley LD, see Stevens JB
Wheeler M, see Dsida RM
White PF, Sá Rêgo Mônica M. The changing role of monitored anesthesia care in the ambulatory setting (letter, reply), 1336
see Fu W
see Sá Rêgo Mônica M.
see Song D
see Tang J
White WD, see Newman MF
Whitten CW, Hill GE, Ivy R, Greilich PE, Lipton JM. Does the duration of cardiopulmonary bypass or aortic cross-clamp, in the absence of blood and/or blood product administration, influence the IL-6 response to cardiac surgery?, 28
see Patel SK
Whitwam JG, see Ma D
Wiedermann CJ, see Manfreda SE
Wiesenfeld-Hallin Z, see Plesan A
Wietlisbach M, see Konrad C
Wilder-Smith OHG, Arendt-Nielsen L, Gäumann D, Tassonyi EA, Rifat KR. Sensory changes and pain after abdominal hysterectomy: a comparison of anesthetic supplementation with fentanyl versus magnesium or ketamine, 95

Wildling E, see Klimscha W
Wilkins LJ, see Groudine SB
Williams E, see Rathmell JP
Williams GD, see Ramamoorthy C
Winnie AP, see Markey JR
Wise JJ, see Hartrick CT
Wong C-S, see Chia Y-Y
Wong JWF, see Riley RH
Wu C-C, see Chou T-C
Wu M-H. Propofol and the supraventricular tachydysrhythmias in children (letter, reply), 914
Wüsten R, see Buerkle H

Xu X-J, see Plesan A
Xue FS, An G, Liao X, Zou Q, Luo LK. The pharmacokinetics of vecuronium in male and female patients, 1322
Liu JH, Liao X, Tong SY, Li L, Zhang RJ, An G, Luo LK. The influence of acute normovolemic hemodilution on the dose-response and time course of action of vecuronium, 861

Yabe M, Nishikawa K, Terai T, Yukioka H, Fujimori M. The effects of intrinsic nitric oxide on cardiac neural regulation in cats, 1194
Yaksh TL, see Jun JH
Yaku H, see Kagawa T
Yamamoto M, Hatano Y, Ogawa K, Iranami H, Tajima T. Halothane and isoflurane attenuate the relaxant response to nonadrenergic and noncholinergic nerve stimulation of isolated canine cerebral arteries, 552
Yamamoto T
see Aoyama K
see Nozaki-Taguchi N
Yamazaki M, Sasaki R, Masuda A, Ito Y. Anesthetic management of complete tracheal disruption using percutaneous cardiopulmonary support system, 998
Yaster M, see Kost-Byerly S
Yen M-H, see Chou T-C
Yernault J-C, see Bardoczky GI
Yokochi A, see Tsujimoto S
Yoshiya I
see Abe K
see Fujino Y
see Shibata M
Yost CS, Hampson AJ, Leonoudakis D, Koblin DD, Bornheim LM, Gray AT. Oleamide potentiates benzodiazepine-sensitive γ-aminobutyric acid receptor activity but does not alter minimum alveolar anesthetic concentration, 1294
Yukioka H, see Yabe M

Zagvazdin Y, Reiner A, Benter IF. How selective is 7-nitroindazole, an inhibitor of neuronal nitric oxide synthase? (letter), 679
Zahn J, see Beilin Y
Zahurak M, see Kost-Byerly S
Zalaquett R, see Lema G
Zamudio S, see Katz JJ
Zarzur E. Intravenous oxytocin in patients undergoing elective cesarean section (letter), 1334
Zbinden AM
see Curatolo M
see Sieber TJ
Zeijlmans PWM, see Groen K
Zenker G, see Trantina AE
Zenzen WJ, see Beilin Y
Zeppenfeldt B, see Groen K
Zetlaoui PJ. Is there a place for meperidine in intravenous regional anesthesia? (letter), 918
Zhang J, Maland L, Hague B, Niu S, Robertson S, Coleman D, Stanley TH, Streisand JB. Buccal absorption of etomidate from a solid formulation in dogs, 1116
see Farnsworth ST
Zhang RJ, see Xue FS
Zheng D, Upton RN, Martinez AM, Grant C, Ludbrook GL. The influence of the bolus injection rate of propofol on its cardiovascular effects and peak blood concentrations in sheep, 1109
Zimpfer M
see Blaicher AM
see Kettner SC
Zornow MH, see Kwon J-Y
Zou Q, see Xue FS
Zucker-Pinchoff B, see Beilin Y

Subject Index

Abdominal surgery, see Surgery
ACE, see Pharmacology, angiotensin-converting enzyme
Acetyl starch, see Pharmacokinetics
Acute normovolemic hemodilution, see Anesthetic techniques
Adenosine triphosphate, see Pharmacology
Adenotonsillectomy, see Surgery
α_2-Adrenergic receptors, see Receptors
Age factors
effects on optimal premedication dose of intramuscular midazolam, 86:1103
elderly
arterial and pulmonary arterial concentrations of bupivacaine enantiomers after epidural injection, 86:812
propofol-induced hypotension, prophylactic effect of ephedrine, female patients, 86:477
surgical population, influence on anesthesia workload, ten-year survey, 86:1165
infants, postoperative, clearance of morphine during intravenous infusion, 86:958
Air release valves, see Equipment
Airway
complications, associated with emergency tracheal reintubation, quality improvement audit, 86:1171
device, emergency, use of Combitube, 86:1137
epithelium, epithelial-derived relaxing factor, volatile anesthetics and bronchodilation, rats, 86:646
management
during resection of high tracheal stenosis, use of LMA and high-frequency jet ventilation, 86:907
intra- and postoperative, patient with total laryngopharyngectomy and cervical esophagectomy (letter), 86:678
trauma, role of anesthesiologists (letter), 86:1146
MRI, infant with micrognathia, 86:964
Airway obstruction, see Complications, airway
Airway occlusion, see Anesthetic techniques
Alae nasi, see Nose
Alfentanil
see Analgesics
see Anesthetics, intravenous
Allergy
anaphylaxis, hydroxyethyl starch-reactive antibody, 86:1123
latex, and anesthesia (letter), 86:919
Allodynia, see Pain, neuropathic
Alveolar macrophages, see Lung
Ambulatory anesthesia, see Anesthesia
Amino acids, excitatory, glutamate, increased hippocampal concentrations during repeated transient global cerebral ischemia, failure of riluzole to attenuate, rabbits, 86:128
γ-Aminobutyric acid, see Receptors
ϵ-Aminocaproic acid, see Antifibrinolytics
Amrinone, see Pharmacology
Amyotrophic lateral sclerosis, see Complications
Analgesics
alfentanil
adjuvant to epidural bupivacaine, management of pain after laparotomy, lack of evidence of spinal action, 86:574
ketamine-alfentanil combinations, vs alfentanil alone, analgesic and cognitive effects after intradermal capsaicin, normal subjects, 86:1250
capsaicin, topical, treatment of intractable pain, preliminary report, 86:579
diclofenac, systemic, combined with small doses of intrathecal morphine, for postoperative pain control after cesarean delivery, 86:538
fentanyl
epidural, vs sufentanil, dose-response function (letter), 86:918
intrathecal, with small-dose dilute bupivacaine, better anesthesia without prolonging recovery (letter), 86:917
premedication, effect on reliability of intravenous lidocaine test dose, 86:1015
small-dose, injection time, optimal, for blunting circulatory responses to tracheal intubation, 86:658
venous retention pretreatment, effect on propofol injection pain, 86:382
gabapentin
intrathecal, effect on hyperalgesia after thermal injury, rats, 86:348
reversal of allodynia produced by administration of anti-GD_2 ganglioside, 86:111
ibuprofen, response, gender differences, healthy human subjects, 86:1257
inositol-triphosphate, postoperative analgesia, patients undergoing cholecystectomy, 86:107
ketorolac
as component of intravenous regional anesthesia with lidocaine, dose-response relationship, 86:791
-associated perioperative renal dysfunction, factors influencing, 86:1090
lornoxicam, PCA with, vs PCA with morphine, patients undergoing lumbar disk surgery, 86:1045
meperidine
intravenous regional anesthesia (letter), 86:918
venous retention pretreatment, effect on propofol injection pain, 86:382
morphine
antinociceptive effect, potentiation, ketamine vs dextromethorphan, rats, 86:825
clearance, postoperative, during intravenous infusion, influence of age and surgery, infants, 86:958
epidural, naloxone-precipitated acute opioid withdrawal syndrome after, 86:544
epidural, side effects, and droperidol, after cesarean section, 86:532
intraarticular, in multimodal analgesic management of pain after ambulatory anterior cruciate ligament repair, 86:374
intraarticular, tourniquet release time and analgesic effects (letter), 86:1150
intrathecal, -induced pruritus, after cesarean section, subhypnotic doses of propofol do not relieve, 86:310
intrathecal, small doses, combined with systemic diclofenac, for postoperative pain control after cesarean delivery, 86:538
PCA, oral vs intravenous, analgesic consumption, 86:1051
PCA, vs PCA with lornoxicam, patients undergoing lumbar disk surgery, 86:1045
peripherally administered, and FK409, interaction, during experimental inflammation, 86:367
sustained-release tablets, prolonged oral treatment, influence on immune function, 86:166

Analgesics—*continued*
venous retention pretreatment, effect on propofol injection pain, 86:382
neostigmine, mediation of central and peripheral analgesia, inflamed knee joint, rats, 86:1027
NSAIDs, epidural, for cancer pain, 86:117
sufentanil
intrathecal, for extracorporeal shock wave lithotripsy, patients with aortic stenosis, 86:943
vs epidural fentanyl, dose-response function (letter), 86:918
Anaphylactoid reactions, see Complications
Anaphylaxis, see Allergy
Anesthesia
ambulatory
effect of timing of ondansetron administration on outcome after ambulatory surgery, 86:274
epinephrine and bupivacaine spinal anesthesia, 86:973
extracorporeal shock wave lithotripsy, general vs epidural anesthesia, 86:1214
fast-track eligibility after ambulatory surgery, 86:267
ondansetron vs droperidol for PONV, 86:731
propofol controlled conscious sedation, 86:967
selective axillary block, 86:746
seven days postoperatively, 86:739
tracheal intubation without muscle relaxant, 86:45
cardiovascular
α_2 agonists on emergence from anesthesia, 86:240
amrinone vs milrinone after cardiac surgery, 86:683
analgesia for extracorporeal shock wave lithotripsy, patients with aortic stenosis, 86:943
ANH and allogeneic transfusion, 86:9
aortic perforation during lumbar laminectomy, 86:493
burst suppression during cardiac surgery, 86:246
cardiac neural regulation, effects of intrinsic nitric oxide, 86:1194
circulatory changes during laparoscopy, 86:482
conjugated estrogen in orthotopic liver transplantation, 86:1183
dysrhythmias and noncardiac surgery, 86:16
effects of thyroid hormone modulation on rat liver injury, 86:1187
ephedrine and propofol hypotension, 86:477
esophageal electrocardiograph and ischemia, 86:34
ethylenediaminetetraacetic acid-dependent pseudothrombocytopenia, 86:1210
etomidate and aortic input impedance, 86:932
gastric mucosal perfusion during CPB, 86:455
halothane and abnormal automaticity, 86:488
heparin activity during liver transplant, 86:691
hepatic changes after CPB and fentanyl, 86:1177
influence of CPB on interleukin 6 response, 86:28
intraoperative torsade de pointes VT and VF during sevoflurane anesthesia, 86:710
intravenous clonidine for liver transplantation anesthesia, 86:468
minimally invasive myocardial revascularization before lung transplantation, 86:939
oxytocin and atrial contractility *in vitro*, 86:40
positive inotropic drugs after valve surgery, 86:461
propofol and ischemic reperfused hearts, 86:252
propofol, desaturation, and hypothermic CPB, 86:926
prostaglandin E_1 and right-to-left shunting, 86:703
renal protection in cardiac surgery, 86:3
right double lumens vs right Univent tubes, 86:696
severe hypercarbia during CPB, complication of CO_2 flooding of surgical field, 86:264
splanchnic oxygen transport and CPB, 86:22
treatment of anesthesia-induced hypotension, patients chronically treated with ACE inhibitors, 86:259
vasopressin for treatment of refractory hypotension after CPB, 86:1207
critical care and trauma
adhesion molecules in traumatic brain injury, 86:759
complete tracheal disruption, anesthetic management, 86:998
dopexamine and ventilation perfusion distribution during anesthesia, 86:314
inhaled prostaglandin E_1 in acute lung injury, 86:753
prolonged inhaled NO, 86:74
propofol vs midazolam, severe trauma patient, 86:1219
dental treatment, conscious sedation with propofol, patient-controlled vs anesthesiologist-controlled, anxious patients, 86:967
electroconvulsive therapy
acute hemodynamic response, modification, effectiveness of oral clonidine, 86:1127
effect on thermal hyperalgesia and mechanical allodynia, rats, 86:584
thermal hyperalgesia and mechanical allodynia, 86:584
neurosurgical
anesthetics and ischemic depolarization, focal ischemia, 86:773
CBF velocity and hemodilution, 86:320
CBF velocity in chronic liver disease, 86:1005
cisatracurium, atracurium, and neurosurgery, 86:123
clonidine and head-holder application, 86:1001
clonidine on pial microcirculation, 86:766
hypothermia and nitroglycerin-induced vasodilation, 86:546
intravenous fluids after closed head trauma, 86:1225
pentobarbital effects on blood-brain barrier disruption, 86:1230
riluzole and glutamate increases during ischemia, 86:128
volatile anesthetics and cerebrovasodilation, 86:522
wavelet analysis and venous air embolism, 86:325
obstetric
amyotrophic lateral sclerosis during pregnancy, 86:542
cesarean section and bupivacaine baricity, 86:989
diaphragmatic hernia during pregnancy, 86:501
effects of epidural isoproterenol, 86:749
epidural morphine and droperidol, 86:532
epidural ropivacaine, labor analgesia, 86:527
epidural test doses, 86:985

epinephrine test dose, limitations (editorial), 86:923
intrathecal morphine and systemic diclofenac, 86:538
mivacurium in postpartum patients, 86:82
mivacurium, magnesium, and neuromuscular block, 86:523
naloxone-precipitated acute opioid withdrawal syndrome after epidural morphine, 86:544
nitroglycerin and uterine relaxation, 86:304
pneumoperitoneum and endobronchial intubation, 86:301
propofol and pruritus, 86:310
sextuplets, cesarean section, successful epidural anesthesia, 86:1236
spinal needles, combined spinal-epidural analgesia, 86:520
sterility of drugs in syringes for long times, 86:994
pediatric
aortic blood flow and laparoscopy, 86:290
consent for pediatric anesthesia research, 86:50
clonidine in pediatric caudal blocks, 86:54
CO_2 embolism during hip arthrography, 86:299
continuous pediatric epidural catheters, 86:712
generalized muscle rigidity and hypercarbia with halothane and isoflurane, 86:297
inspiratory collapse of alae nasi causing variable extrathoracic airway obstruction, 86:294
LMAs for upper respiratory tract infection, 86:703
management of congenital lobar emphysema with endobronchial intubation and controlled ventilation, 86:71
milrinone after heart surgery, 86:283
mivacurium after atracurium or cisatracurium, 86:62
morphine clearance, infants, 86:958
MRI of airway, infant with micrognathia, 86:964
mucormycosis supraglottitis on anesthetic induction, immunocompromised host, 86:729
oral transmucosal fentanyl for tonsillectomy, 86:66
propofol and newborn cardiac function, 86:717
sedation with chloral hydrate and N_2O, 86:724
test dose efficacy, 86:952
tracheal intubation for pyloromyotomy, 86:945
video-assisted thoracoscopic surgery, modified endotracheal tube, 86:1212
vomiting after adenotonsillectomy, 86:496
regional anesthesia and pain management
addition of CO_2 and HCO_3 to lidocaine, 86:341
analgesia by inositol-triphosphate, 86:107
anesthesia and severe hypercapnia, 86:591
bupivacaine and mepivacaine enantiomers, 86:361
bupivacaine enantiomer pharmacokinetics, 86:805
clonidine and epidural test dose, 86:1010
dexmedetomidine and neuropathic pain, 86:355
diaphragmatic paralysis complicating alcohol splanchnic nerve block, 86:845
effects of cervical epidural on breathing, 86:1033
enantiomers of bupivacaine-lung uptake, 86:812
epidural alfentanil for postoperative analgesia, 86:574
epidural anesthesia and gastrointestinal motility, 86:837
epidural NSAIDs for cancer pain, 86:117
epidural sufentanil administered after dural puncture, 86:794
epinephrine and spinal anesthesia, 86:1018
experimental pain in healthy human subjects, 86:1257
gabapentin for neuropathic pain, 86:111
intrathecal ketamine isomers, rats, 86:561
intrathecal magnesium and morphine, 86:830
ketamine-alfentanil combinations after intradermal capsaicin, 86:1250
ketamine delays neuropathic hyperalgesia, 86:557
ketamine in PCEA regimen, 86:1245
ketorolac dose-response in intravenous regional anesthesia, 86:791
lidocaine-related neurologic syndrome, 86:1023
large-dose capsaicin and intractable pain, 86:579
lornoxicam and morphine in acute pain, 86:1045
lumbar sympathetic ganglion block, patient with painful legs and moving toes syndrome, 86:1056
N-methyl-D-aspartate antagonists and morphine, 86:825
multimodal analgesia for anterior cruciate ligament repair, 86:374
nerve toxicity with lidocaine, 86:569
new indicator of somatosensory blockade, 86:336
NO and peripheral morphine analgesia, 86:367
onset of axillary blockade and alkalinization, 86:566
opioid peripheral analgesic effects, 86:382
pain during intradermal anesthesia, 86:379
patient-controlled analgesia, intravenous vs oral, 86:1051
patient-controlled regional analgesia at home, 86:86
posterior subcapsular cataract formation and lumbar epidural corticosteroid injections, 86:1054
postoperative analgesia with epidural verapamil, 86:786
postoperative pain and delirium, 86:781
respiratory, motor, and sensory function after supraclavicular block, 86:1239
sedation and intravenous lidocaine test dose, 86:1015
sensory block and thoracic epidural, 86:332
sensory change after surgery, 86:95
spinal and peripheral analgesic action of neostigmine, 86:1027
spinal gabapentin antihyperalgesia, 86:348
sympathetic blockade and pain from veins, 86:588
systemic toxicity of levobupivacaine, 86:797
three-in-one vs fascia iliaca compartment block, 86:1039
unsuccessful resuscitation under hypotensive epidural anesthesia, elective hip arthroplasty, 86:847
vasovagal cardiac arrest during epidural catheter insertion, 86:1263

Anesthesia—*continued*
verapamil, nimodipine, and bupivacaine, 86:818
vision loss after epidural injection, 86:119
visual analog scale in postoperative period, 86:102
Vo_2 and Vco_2 after tourniquet release, 86:90
Anesthesiologists
formal events, accessories (letter), 86:219
role, management of trauma airway (letter), 86:1146
use of intraoperative resources, effect of drug and supply cost feedback system, 86:510
Anesthetics, gases, nitrous oxide
additive sedative and respiratory effects, chloral hydrate sedation, 86:724
effect on chest wall function, humans and dogs, 86:1058
effect on nociceptive responses, 86:629
Anesthetics, intravenous
alfentanil, recovery from anesthesia, clinical implication of context-sensitive half-times, 86:184
clonidine, effects on catecholamine activation, vasomotor center, emergence from anesthesia, 86:240
etomidate
buccal absorption from solid formulation, dogs, 86:1116
adverse alteration of determinants of left ventricular afterload, dogs with dilated cardiomyopathy, 86:932
fentanyl
influences on hepatic circulation and oxygen metabolism, beagles, 86:1177
interaction with propofol, 86:872
vs magnesium or ketamine, anesthetic supplementation, sensory changes and pain after abdominal hysterectomy, 96:95
idazoxan, effects on catecholamine activation, vasomotor center, emergence from anesthesia, 86:240
ketamine
-alfentanil combinations, vs ketamine alone, analgesic and cognitive effects after intradermal capsaicin, normal subjects, 86:1250
effect on production and release of endothelin 1 from cultured bovine endothelial cells, 86:1098
intrathecal, antinociceptive effect on carrageenan hyperalgesia, rats, 86:561
intrathecal, preemptive, effect on mechanical hyperalgesia, rats, 86:557
multimodal PCEA regimen, 86:1245
vs dextromethorphan, potentiation of antinociceptive effect of morphine, rats, 86:825
vs fentanyl, anesthetic supplementation, sensory changes and pain after abdominal hysterectomy, 96:95
vs ketamine-alfentanil combinations, analgesic and cognitive effects after intradermal capsaicin, normal subjects, 86:1250
vs midazolam, inhibitory effects on human neutrophil function, 86:159
vs thiopental, inhibitory effects on human neutrophil function, 86:159
vs volatile anesthetics, alterations in course of acid-induced lung injury after general anesthesia, rats, 86:141
lidocaine, effects on postoperative pain, hospital length-of-stay, and return of bowel function, patients undergoing radical retropubic prostatectomy, 86:235
midazolam
effect on interleukin-8 from human polymorphonuclear leukocytes, 86:1289
effects in propofol-induced anesthesia (letter), 86:1148
intramuscular, optimal premedication dose, effects of age and gender, 86:1103
vs ketamine, inhibitory effects on human neutrophil function, 86:159
vs propofol, safety and efficacy for sedating severe trauma patient, 86:1219
vs thiopental, inhibitory effects on human neutrophil function, 86:159
mivazerol, effects on catecholamine activation, vasomotor center, emergence from anesthesia, 86:240
pentobarbital
effects on blood-brain barrier disruption caused by intracarotid injection of hyperosmolar mannitol, rats, 86:1230
reduction of frequency of transient ischemic depolarization during focal ischemia, rats, 86:773
propofol
and remifentanil, for tracheal intubation without muscle relaxants, ambulatory surgery patients, 86:45
and supraventricular tachydysrhythmias, children (letter), 86:914
and thiopental, 1:1 volume mixture, chemical stability, 86:422
bolus injection rate, influence on cardiovascular effects and peak blood concentrations, sheep, 86:1109
brain concentrations, effect of rate of administration, sheep, 86:1301
concentration required for endotracheal intubation with laryngoscope or fiberscope, and interaction with fentanyl, 86:872
conscious sedation, patient-controlled vs anesthesiologist-controlled, for dental treatment in anxious patients, 86:967
effect on interleukin-8 from human polymorphonuclear leukocytes, 86:1289
effects on CBF velocity and cerebral oxygen extraction during CPB, 86:1201
effects on functional and metabolic recovery, ischemic reperfused isolated rat hearts, 86:252
failure to ameliorate cerebral venous oxyhemoglobin desaturation during hypothermic CPB, 86:926
-induced anesthesia, effects of midazolam (letter), 86:1148
-induced hypotension, prophylactic effect of ephedrine, elderly female patients, 86:477
infusion, unaltered left ventricular systolic diastolic function, newborn swine, 86:717
injection pain, opioid peripheral analgesic effects, 86:382
recovery from anesthesia, clinical implication of context-sensitive half-times, 86:184
sedation, production of dose-dependent suppression of lidocaine-induced seizures, rats, 86:652
subhypnotic doses, failure to relieve intrathecal-induced pruritus after cesarean section, 86:310
vs desflurane, economic considerations, 86:504
vs desflurane, fast-track eligibility after ambulatory anesthesia, 86:267
vs isoflurane, economic considerations, 86:504
vs midazolam, safety and efficacy for sedating severe trauma patient, 86:1219

vs sevoflurane, economic considerations, 86:504
vs sevoflurane, fast-track eligibility after ambulatory anesthesia, 86:267
remifentanil
and propofol, for tracheal intubation without muscle relaxants, ambulatory surgery patients, 86:45
tolerance, rapid development, humans, 86:1307
sufentanil
CSF concentrations, after administration of epidural sufentanil, effect of prior dural puncture, sheep, 86:794
ocular transmucosal absorption and toxicity, dogs, 86:138
recovery from anesthesia, clinical implication of context-sensitive half-times, 86:184
thiopental
and propofol, 1:1 volume mixture, chemical stability, 86:422
prepared solutions, chemical stability and sterility, 86:208
vs isoflurane, pharmacologic electroencephalographic suppression during CPB, 86:246
vs ketamine, inhibitory effects on human neutrophil function, 86:159
vs midazolam, inhibitory effects on human neutrophil function, 86:159

Anesthetics, local
bupivacaine
cervical epidural, effects on pulmonary function, conscious patients, 86:1033
clonidine/bupivacaine combination, efficacy and safety, caudal blockade for pediatric hernia repair, 86:54
enantiomers, arterial and pulmonary arterial concentrations after epidural injection, elderly patients, 86:812
enantiomers, pharmacokinetics, after epidural administration of racemates, 86:361
enantiomers, systemic and regional pharmacokinetics, sheep, 86:805
epidural, alfentanil as adjuvant to, management of pain after laparotomy, lack of evidence of spinal action, 86:574
inappropriately large dose, as possible cause of life-threatening upper airway obstruction after glossopharyngeal nerve block (letter), 86:678
-induced cardiotoxicity, effects of verapamil and nimodipine, *in vivo* and *in vitro* study, rats, 86:818
intraarticular, in multimodal analgesic management of pain after ambulatory anterior cruciate ligament repair, 86:374
intravenous, cardiovascular and central nervous system effects, sheep, 86:797
0.125%, vs ropivacaine 0.125%, labor analgesia using patient-controlled epidural infusion, 86:527
small-dose dilute, intrathecal fentanyl with, better anesthesia without prolonging recovery (letter), 86:917
small-dose hyperbaric, effect of epinephrine, clinical implications for ambulatory surgery, 86:973
chloroprocaine, use in intravenous regional anesthesia (letter, reply), 86:1334
EMLA cream, improvement of pulse oximetry signal (letter), 86:915
levobupivacaine
intravenous, cardiovascular and central nervous system effects, sheep, 86:797
systemic and regional pharmacokinetics, sheep, 86:805
lidocaine
alkalinization, failure to hasten onset of axillary brachial plexus block, 86:566
-induced seizures, dose-dependent suppression by propofol sedation, rats, 86:652
intravenous regional anesthesia, ketorolac as component, dose-response relationship, 86:791
neurotoxicity, manifestations, graded, irreversible changes in giant axons, crayfish, 86:569
plus sodium bicarbonate, effect on pain during intradermal injection, 86:379
plus sodium bicarbonate, enhancement of depth of epidural blockade, 86:341
spinal anesthesia, duration, thoracic and lumbosacral dermatomes, effect of varied doses of epinephrine, 86:1018
spinal anesthesia, transient neurologic syndrome in 1045 patients, 86:1023
test dose, reliability, effect of premedication with fentanyl and midazolam, 86:1015
test dose, thoracic epidural administration, three different levels, extension of sensory blockade, 86:322
venous retention pretreatment, effect on propofol injection pain, 86:382
mepivacaine, enantiomers, pharmacokinetics, after epidural administration of racemates, 86:361
motor and sensory block duration, 86:746
postoperative PCA, home environment, 86:86
ropivacaine, 0.125%, vs 0.125% bupivacaine, labor analgesia using patient-controlled epidural infusion, 86:527
spread, three-in-one vs fascia iliaca compartment block, 86:1039
toxicity, treatment, midazolam, flumazenil administered to antagonize, triggering of opisthotonos (letter), 86:677

Anesthetics, volatile
desflurane
bronchodilatory effect, epithelial dependence, distal bronchi, rats, 86:646
case of suspected malignant hyperthermia (letter), 86:916
"cross-over" from isoflurane toward end of anesthesia, recovery after, rats breathing from nonrebreathing system, 86:198
effect on nociceptive responses, 86:629
nonimmobilizers, ventilatory effects, swine, 86:173
MAC, failure of oleamide to alter, 86:1294
malignant hyperthermia associated with, humans, 86:1328
suspected malignant hyperthermia episode during anesthesia (letter), 86:449
vs isoflurane, economic considerations, 86:504
vs propofol, economic considerations, 86:504
vs propofol, fast-track eligibility after ambulatory anesthesia, 86:267
vs sevoflurane, economic considerations, 86:504
vs sevoflurane, effect of anesthetic duration on kinetic and recovery characteristics, volunteers, 86:414
vs sevoflurane, fast-track eligibility after ambulatory anesthesia, 86:267

Anesthetics, volatile—*continued*
enflurane
inhibition of *Pseudomonas aeruginosa* growth, culture conditions reproducing alveolar environment, 86:1075
vs ketamine, alterations in course of acid-induced lung injury after general anesthesia, rats, 86:141
halothane
anesthesia, effect of N_2O on chest wall function, humans and dogs, 86:1058
anesthesia, inhibitory influence of pulmonary vagal afferents on respiratory distress induced by airway occlusion, cats, 86:398
circuit absorption, 86:1070
depth of anesthesia, airway occlusion, new concept of MAC-airway occlusion response, cats, 86:191
effect on fatigue-induced changes in diaphragmatic contractility, hamsters, 86:392
effects on abnormal automaticity, cardiac Purkinje fibers, dogs, 86:488
effects on single human neuronal L-type calcium channels, 86:885
inhibition of Pseudomonas aeruginosa growth, culture conditions reproducing alveolar environment, 86:1075
unusual response, generalized muscle rigidity and hypercarbia, 86:297
vs isoflurane, attenuation of relaxant response to nonadrenergic and noncholinergic nerve stimulation, dogs, 86:552
vs ketamine, alterations in course of acid-induced lung injury after general anesthesia, rats, 86:141
isoflurane
anesthesia, ventilatory responses to acute and sustained hypoxia, 86:403
-anesthetized adults, combination of epinephrine and isoproterenol as simulated epidural test dose, 86:1312
awakening concentration, effect of oral clonidine premedication, 86:410
circuit absorption, 86:1070
"cross-over" to desflurane toward end of anesthesia, recovery after, rats breathing from nonrebreathing system, 86:198
depth of anesthesia, new concept of MAC-airway occlusion response, cats, 86:191
general anesthesia, awareness with recall, incidence and risk factors, 86:1084
inhibition of *Pseudomonas aeruginosa* growth, culture conditions reproducing alveolar environment, 86:1075
prolonged administration, patients with severe renal dysfunction, 86:440
reduction of frequency of transient ischemic depolarization during focal ischemia, rats, 86:773
unusual response, generalized muscle rigidity and hypercarbia, 86:297
vs desflurane, economic considerations, 86:504
vs halothane, attenuation of relaxant response to nonadrenergic and noncholinergic nerve stimulation, isolated cerebral arteries, dogs, 86:552
vs ketamine, alterations in course of acid-induced lung injury after general anesthesia, rats, 86:141
vs propofol, economic considerations, 86:504
vs sevoflurane, arterial oxygenation and shunt fraction during one-lung ventilation, 86:1266
vs sevoflurane, economic considerations, 86:504
vs thiopental, pharmacologic EEG suppression during CPB, 86:246
sevoflurane
anesthesia, intraoperative torsade de pointes VT and VF, 86:701
-anesthetized children, simulation of epidural test dose with intravenous epinephrine, 86:952
bronchodilatory effect, epithelial dependence, distal bronchi, rats, 86:646
circuit absorption, 86:1070
depth of anesthesia, new concept of MAC-airway occlusion response, cats, 86:191
effect on fatigue-induced changes in diaphragmatic contractility, hamsters, 86:392
effect on spontaneous sympathetic activity, Aδ and C somatosympathetic reflexes, and associated hemodynamic changes, dogs, 86:1079
low-flow delivery, four hours of 1.25 MAC, absence of renal and hepatic toxicity, volunteers, 86:662
requirements for MAC and MAC-awake, effect of adenosine triphosphate, 86:179
vs desflurane, economic considerations, 86:504
vs desflurane, effect of anesthetic duration on kinetic and recovery characteristics, volunteers, 86:414
vs desflurane, fast-track eligibility after ambulatory anesthesia, 86:267
vs isoflurane, arterial oxygenation and shunt fraction during one-lung ventilation, 86:1266
vs isoflurane, economic considerations, 86:504
vs propofol, economic considerations, 86:504
vs propofol, fast-track eligibility after ambulatory anesthesia, 86:267
Anesthetic techniques
acute normovolemic hemodilution
and reduction of perioperative allogeneic transfusion, meta-analysis, 86:9
influence on dose-response and time course of action of vecuronium, 86:861
airway occlusion
-induced respiratory distress, inhibitory influence of pulmonary vagal afferents, halothane-anesthetized cats, 86:398
new concept of MAC-airway occlusion response, cats, 86:191
pressure, effects of cervical epidural on breathing, 86:1033
brachial plexus block
axillary, onset, failure of alkalinization of lidocaine to hasten, 86:566
axillary, selective, use for outpatient hand surgery, 86:746
"midhumeral" approach (letter), 86:447
caudal blockade, clonidine/bupivacaine combination, efficacy and safety, pediatric hernia repair, 86:54
cervical epidural, with bupivacaine, effects on pulmonary function, conscious patients, 86:1033
combined spinal-epidural
and incidence of paresthesia during epidural catheter placement (letter), 86:448
four spinal needles, *in vivo* evaluation, 86:520
conscious sedation
with chloral hydrate and N_2O, 86:724
with propofol, patient-controlled vs anesthesiologist-controlled, for

dental treatment in anxious patients, 86:967
continuous spinal anesthesia
safety (letter), 86:218
successful, patient with preoperative baseline hypercapnia exceeding 100 mm Hg, 86:591
vs single-injection spinal anesthesia, complications (letter), 86:214
epidural
administration, bupivacaine and mepivacaine racemates, pharmacokinetics of enantiomers, 86:361
and gastrointestinal motility, 86:837
anesthesia, cesarean section, sextuplets, 86:1236
blockade, and perioperative low molecular weight heparin (editorial), 86:1153
blockade, depth, enhanced by adding sodium bicarbonate to lidocaine, 86:341
bupivacaine, arterial and pulmonary arterial concentrations of bupivacaine enantiomers after injection, elderly, 86:812
catheter insertion, vasovagal cardiac arrest during, before administration of epidural medication, 86:1263
catheters, continuous, bacterial colonization and infection rate, children, 86:712
clonidine, spinal cord accumulation, effects of epidural epinephrine, rabbits (letter), 86:1145
morphine, naloxone-precipitated acute opioid withdrawal syndrome after, 86:544
morphine, side effects, and droperidol, after cesarean section, 86:532
NSAIDs for cancer pain, 86:117
patients breathing spontaneously, changes in metabolic variables after tourniquet release, 86:90
quality and duration, effects of adding isoproterenol to 0.125% bupivacaine, laboring parturients, 86:749
steroid injection, lumbar, vision loss after, 86:119
test dose, epinephrine, obstetrics, limitations (editorial), 86:923
test dose, epinephrine, optimal, for epidural injection with lidocaine solution, awake patients premedicated with oral clonidine, 86:1010
test dose, epinephrine, prospective clinical study, laboring women, 86:985
test dose, simulated, combination of epinephrine and isoproterenol, isoflurane-anesthetized adults, 86:1312
test dose, simulation, with intravenous epinephrine, sevoflurane-anesthetized children, 86:952
verapamil, reduction of analgesic consumption after lower abdominal surgery, 86:786
vs general, for outpatient extracorporeal shock wave lithotripsy, 86:1214
fascia iliaca compartment block, vs three-in-one block, clinical and radiographic analysis, adults, 86:1039
fiberscopy, required propofol concentration, 86:872
general
awareness with recall, incidence and risk factors, 86:1084
consensus (letter), 86:446
heart and lung sound monitoring, effect of esophageal stethoscope placement depth, 86:1276
vs epidural, for outpatient extracorporeal shock wave lithotripsy, 86:1214
glossopharyngeal nerve block, life-threatening upper airway obstruction after, inappropriately large dose of bupivacaine as possible cause (letter), 86:678
hypotensive epidural anesthesia, unsuccessful resuscitation during elective hip arthroplasty, 86:847
injection rate, propofol, influence on cardiovascular effects and peak blood concentrations, sheep, 86:1109
interscalene block, for outpatient shoulder arthroscopy (letter), 86:217
intravenous regional
use of chloroprocaine (letter, reply), 86:1334
use of meperidine (letter), 86:918
with lidocaine, ketorolac as component, dose-response relationship, 86:791
laryngoscopy, required propofol concentration, 86:872
low-flow, four hours of 1.25 MAC sevoflurane, absence of renal and hepatic toxicity, volunteers, 86:662
one-lung ventilation
arterial oxygenation and shunt fraction, isoflurane vs sevoflurane, 86:1266
positioning left-sided double-lumen tube without aid of fiberoptic bronchoscopy, 86:1271
positive end-expiratory pressure, intrinsic, and interrupted expiratory flow on automatically constructed flow-volume curves, 86:880
right-sided double-lumen tube vs single-lumen tube with right-sided enclosed bronchial blocker (Univent), 86:696
oral mucosal administration, buccal absorption of etomidate from solid formulation, dogs, 86:1116
patient-controlled analgesia
intravenous vs oral, analgesic consumption, 86:1051
local anesthetic administration, home environment, 86:86
with lornoxicam, vs PCA with morphine, patients undergoing lumbar disk surgery, 86:1045
patient-controlled epidural analgesia
after abdominothoracic esophagectomy, 86:228
for labor analgesia, 0.125% ropivacaine vs 0.125% bupivacaine, 86:527
multimodal regimen, adding ketamine, effect on postoperative pain and analgesic consumption, 86:1245
percutaneous cardiopulmonary support system, for anesthetic management of complete tracheal disruption, 86:998
preemptive, ketamine delays neuropathic hyperalgesia, rats, 86:557
regional, level, assessment, objective indicator, use of skin vasomotor reflex, 86:336
spinal
analgesic action of neostigmine, inflamed knee joint, rats, 86:1027
blockade, and perioperative low molecular weight heparin (editorial), 86:1153
lidocaine, duration, thoracic and lumbosacral dermatomes, effect of varied doses of epinephrine, 86:1018
lidocaine, 3%, transient neurologic syndrome in 1,045 patients, 86:1023

Anesthetic techniques—*continued*
morphine, co-infusion of magnesium sulfate, antinociceptive potentiation and attenuation of tolerance, rats, 86:830
morphine, small doses, combined with systemic diclofenac, postoperative pain control after cesarean delivery, 86:538
small-dose hyperbaric bupivacaine, effect of epinephrine, clinical implications for ambulatory surgery, 86:973
small-dose hyperbaric bupivacaine, vs plain bupivacaine, for cesarean section, 86:989
splanchnic nerve block, alcohol, complicated by diaphragmatic paralysis, 86:845
supraclavicular nerve block
respiratory, motor, and sensory function after, quantitative analysis, 86:1239
sternocleidomastoid muscle, surface landmark (letter), 86:1149
suprascapular block, shoulder arthroscopy (letter), 86:217
sympathetic block
failure to reduce acute vascular pain, humans, 86:588
lumbar sympathetic ganglion block, patient with painful legs and moving toes syndrome, 86:1056
thoracic epidural, test dose of lidocaine, three different levels, extension of sensory blockade, 86:332
three-in-one block
femoral, ultrasonic performance (letter), 86:1147
vs fascia iliaca compartment block, clinical and radiographic analysis, adults, 86:1039
total intravenous anesthesia, patient with malignant systemic mastocytosis, 86:442
Angiotensin-converting enzyme inhibitors, see Pharmacology
Angiotensin II, see Pharmacology
Antagonists, opioid, naloxone
effect on chemotaxis of human peripheral blood B lymphocytes by β-endorphin, 86:670
-precipitated acute opioid withdrawal syndrome after epidural morphine, 86:544
Antibiotics, gentamicin, ketorolac plus, production of perioperative renal dysfunction, 86:1090
Antibodies
anti-GD_2, administration, allodynia produced by, reversal by gabapentin, 86:111
formation, due to hydroxyethyl starch, humans, incidence and clinical relevance, 86:1123
Antiemetics
dimenhydrinate, vs ondansetron, effectiveness in prevention of vomiting after adenotonsillectomy, children, 86:496
droperidol, and granisetron, combination, prevention of PONV, 86:613
granisetron, and droperidol, combination, prevention of PONV, 86:613
ondansetron
administration, timing, effect on efficacy, cost-effectiveness, and cost-benefit as prophylactic antiemetic, 86:274
vs dimenhydrinate, effectiveness in prevention of vomiting after adenotonsillectomy, children, 86:496
tropisetron, for treating established PONV, randomized, double-blind, placebo-controlled study, 86:617
Antifibrinolytics, ε-aminocaproic acid, plasma levels, measurement during CPB (letter), 86:680
Antinociception, see Pain
Aorta, blood flow, monitoring, continuous esophageal echo-Doppler, assessment of hemodynamic effects of pneumoperitoneum during laparoscopic surgery, healthy infants, 86:283
Aortic perforation, see Complications
Aortic stenosis, see Heart
L-Arginine transport, see Lung, alveolar macrophages
Arrhythmias, see Heart
Arteries
cerebral, relaxant response to nonadrenergic and noncholinergic nerve stimulation, effects of halothane vs isoflurane, dogs, 86:552
pial, nitroglycerin-induced vasodilation, effect of mild hypothermia, cats, 86:546
pulmonary, concentrations of bupivacaine enantiomers after epidural injection, elderly patients, 86:812
Arthroplasty
see Surgery, hip
see Surgery, knee
Arthroscopic surgery, see Surgery
Aspiration of gastric contents, see Complications
Atracurium, see Neuromuscular relaxants
Atrial septal defect, see Heart
Atrial trabeculae, see Heart
Augustine Guide, see Equipment
Awareness, see Complications
Axillary block, see Anesthetic techniques, brachial plexus block

Bacterial growth, see Infection
Barbiturates, therapy, and severe hypokalemia (letter), 86:214
Benzodiazepines, -sensitive γ-aminobutyric acid receptor activity, potentiation by oleamide, 86:1294
Blood
-brain barrier, disruption caused by intracarotid injection of hyperosmolar mannitol, effects of pentobarbital, rats, 86:1230
coagulation
effect of hydroxyethyl starch, 86:1318
endogenous heparin-like substances, cause of significant coagulation impairment, patients undergoing orthotopic liver transplantation, 86:691
hypercoagulable fibrinolysis, postoperative, failure of neuroendocrine stress hormones to recreate, 86:640
management, during orthotopic liver transplantation, use of conjugated estrogen, 86:1183
monitoring, thromboelastography, parturient with immune thrombocytopenic purpura (letter), 86:675
hemodilution
effect on allogeneic blood requirements, patients undergoing hip or knee arthroplasty, 86:387
effect on CBF velocity, anesthetized patients, 86:320
peripheral, B lymphocytes, chemotaxis by β-endorphin, reversibility by naloxone, humans, 86:670
plasma substitutes
acetyl starch, vs hydroxyethyl starch, pharmacokinetics, patients undergoing elective surgery, 86:856
hydroxyethyl starch, antibodies, humans, incidence and clinical relevance, 86:1123
hydroxyethyl starch, effect on platelet aggregation in vitro, 86:1318

hydroxyethyl starch, 6% solution, assessment of safety and tolerance, randomized, controlled epidemiology study, 86:850
hydroxyethyl starch, vs acetyl starch, pharmacokinetics, patients undergoing elective surgery, 86:856
platelets, aggregation, *in vitro*, effect of hydroxyethyl starch, 86:1318
red cells, scavenging, effect on allogeneic blood requirements, patients undergoing hip or knee arthroplasty, 86:387
transfusion
absence, duration of CPB or aortic cross-clamp, influence on IL-6 response to cardiac surgery, 86:28
allogeneic, perioperative, reduction, and acute normovolemic transfusion, meta-analysis, 86:9
allogeneic, requirements, effects of red-cell scavenging, hemodilution, and active warming, hip or knee arthroplasty, 86:387
requirements, during orthotopic liver transplantation, reducing, efficacy of conjugated estrogen, 86:1183

Blood flow
see Aorta
see Brain
see Liver

Blood pressure
drug effects, clonidine and epidural test dose, 86:1010
effects of cisatracurium vs atracurium, neurosurgical patients, 86:123
hypotension
anesthesia-induced, patients chronically treated with ACE inhibitors, treatment, use of angiotensin II, 86:259
caused by propofol, influence of bolus injection rate, sheep, 86:1109
effect of sevoflurane, dogs, 86:1079

Book and multimedia reviews
Acute and Chronic Pain, Vol. 25, No. 2 of International Anesthesiology Clinics, 86:682
Airway Management in Pediatric Anesthesia. Vol. 35, No. 3 of International Anesthesiology Clinics, 86:921
Anesthesia: A Comprehensive Review, 86:221
Anesthesia and Perioperative Care of the Combat Casualty, 86:222
Anesthesia and Uncommon Diseases. 4th ed., 86:1151
Anesthesia Education at Emory 1958–1985, 86:452
Anesthesia for Cardiac Surgery, 86:1340
Anesthesia for the Patient with Neurologic Disease. Anesthesiology Clinics of North America. Vol 15, No. 3, 86:453
Critical Care Medicine in Transition: The Anesthesiologist's Role. Vol. 15, No 4 of Anesthesiology Clinics of North America, 86:1341
Direct Laryngoscopy Video System, 86:921
Handbook of Mechanical Ventilatory Support, 2nd ed., 86:222
Handbook of Palliative Care in Cancer, 86:221
The ICU Book. 2nd ed., 86:1151
The Laryngeal Mask Airway: A Review and Practical Guide, 86:452
Management of the Elderly Surgical Patient. Vol. 9, No. 4 of Problems in Anesthesia, 86:1152
Neuroprotection: A Society of Cardiovascular Anesthesiologists Monograph, 86:921
Pain: Nociceptive and Neuropathic Mechanisms with Clinical Correlates. Vol. 15, No. 2 of Anesthesiology. Clinics of North America, 86:221
Pediatric Cardiac Anesthesia, 86:1341
Perioperative Care: Anesthesia, Medicine, and Surgery, 86:922
Principles and Practice of Anesthesiology. 2nd Ed., 86:1340

Bowel function, see Gastrointestinal tract

Brachial plexus block, Anesthetic techniques

Brain
adhesion molecules, soluble, concentrations, in systemic and jugular venous serum, and injury severity and outcome after traumatic brain injury, relationship, 86:759
blood-brain barrier, disruption caused by intracarotid injection of hyperosmolar mannitol, effects of pentobarbital, rats, 86:1230
blood flow
cerebral arteries, relaxant response to nonadrenergic and noncholinergic nerve stimulation, effects of halothane vs isoflurane, dogs, 86:552
velocity, during CPB, effects of propofol, 86:1201
velocity, effect of hemodilution, 86:320
velocity, patients with subclinical portal-systemic encephalopathy, 86:1005
cerebral perfusion pressure, effects of cisatracurium vs atracurium, neurosurgical patients, 86:123
cerebral venous oxyhemoglobin desaturation, during hypothermic CPB, failure of propofol to ameliorate, 86:926
circulation, pial arteriolar diameter, nitroglycerin-induced vasodilation, effect of mild hypothermia, cats, 86:546
concentrations of propofol, effect of rate of administration, sheep, 86:1301
death, organ donors, effects of thyroid hormone modulation, 86:1187
edema, two hours after closed head trauma, effects of intravenous fluids, rats, 86:1225
intracranial pressure, effects of cisatracurium vs atracurium, neurosurgical patients, 86:123
ischemia
focal, frequency of transient ischemic depolarization, effects of isoflurane and pentobarbital, rats, 86:773
transient global, increased hippocampal glutamate concentrations during, failure of riluzole to attenuate, rabbits, 86:128
lipids, oleamide, potentiation of benzodiazepine-sensitive γ-aminobutyric acid receptor activity, 86:1294
metabolism, pharmacologic EEG suppression during CPB, thiopental vs isoflurane, 86:246
oxygen extraction, during CPB, effects of propofol, 86:1201
pial microcirculation, vascular effects of topical and intravenous clonidine, dogs, 86:766
regional cerebral oxygenation, differential effects of prostaglandin E_1 and nitroglycerin (letter, reply), 86:1338, 1339
traumatic injury
closed head trauma, edema after, effects of intravenous fluids, rats, 86:1225
severity and outcome, and soluble adhesion molecule concentrations in systemic and jugular venous serum, relationship, 86:759
vasomotor center, catecholamine activation on emergence from

Brain—*continued*
anesthesia, effects of α_2 agonists, 86:240
Bronchodilation, see Lung

Calcium, see Ions
Cancer
see Complications, laryngeal carcinoma
see Pain
Capnography, see Measurement techniques
Capsaicin, see Analgesics
Carbon dioxide
absorbents, degradation of volatile anesthetics, 86:1070
added to lidocaine, lack of effect on depth of epidural blockade, 86:341
elimination, changes after tourniquet release, patients breathing spontaneously under epidural anesthesia, 86:90
embolism, during hip arthroplasty, infant, 86:299
flooding of surgical field during CPB, cause of severe hypercarbia, 86:264
gradient, arterial to end-tidal, increases with uncorrected but not with temperature-corrected $Paco_2$ determination, mild to moderate hypothermia, 86:1131
hypercapnia, preoperative baseline, exceeding 100 mm Hg, successful continuous spinal anesthetic, 86:591
hypercarbia, and generalized muscle rigidity, with halothane and isoflurane, 86:297
insufflation, peritoneal, pneumoperitoneum
effect on intraocular pressure, α-chymotrypsin-induced glaucoma, rabbits, 86:1283
hemodynamic effects, assessment by continuous esophageal aortic blood flow echo-Doppler, laparoscopic surgery, healthy infants, 86:290
-induced left ventricular loading modifications, time course echocardiographic study, 86:482
risk factor for endobronchial intubation, laparoscopic gynecologic surgery, 86:301
noninvasive monitoring, end-tidal vs transcutaneous CO_2 (letter), 86:675
reactivity, patients with liver disease, 86:1005
tensions, during anesthesia in prone position, 86:668
Cardiac arrest, see Complications
Cardiopulmonary bypass, see Surgery
Cardiotoxicity, see Toxicity
Cardiovascular anesthesia, see Anesthesia
Cataract formation, see Complications
Catecholamines, see Sympathetic nervous system
Catheters, see Equipment
Caudal blockade, see Anesthetic techniques
CBF, see Brain, blood flow
Central core disease, see Complications
Central venous pressure, see Monitoring
Cerebellar hemangioblastoma excision, see Surgery
Cerebral arteries, see Arteries
Cerebrospinal fluid, concentrations of sufentanil after epidural administration, effect of prior dural puncture, sheep, 86:794
Cervical epidural, see Anesthetic techniques
Cesarean section, see Surgery
Chest wall, see Muscle
Chloral hydrate, see Hypnotics
Chlorobutanol, see Preservatives
Chloroprocaine, see Anesthetics, local
Cholecystectomy, see Surgery
Cholinesterase, see Enzymes
Chronic obstructive pulmonary disease, see Lung
Cisatracurium, see Neuromuscular relaxants
Clinical pathway, perioperative, for knee replacement surgery, effect on hospital costs, 86:978
Clonidine
see Anesthetics, intravenous
see Pharmacology
see Premedication
see Sympathetic nervous system
Coagulation, see Blood
Collapsed anterior chamber, see Complications
Combined spinal-epidural, see Anesthetic techniques
Combitube, see Equipment
Complete tracheal disruption, see Complications
Complications
airway
associated with emergency tracheal reintubation, quality improvement audit, 86:1171
obstruction, extrathoracic, caused by inspiratory collapse of alae nasi, 86:294
obstruction, failure of LMA, undiagnosed laryngeal carcinoma, 86:438
obstruction, life-threatening, after glossopharyngeal nerve block, inappropriately large dose of bupivacaine as possible cause (letter), 86:678
amyotrophic lateral sclerosis, presenting during pregnancy, 86:542
anaphylactoid reactions, after administration of 6% hydroxyethyl starch solution, randomized, controlled epidemiology study, 86:850
aortic perforation, during lumbar laminectomy, 86:493
aspiration of gastric contents, acid-induced lung injury, effects of ketamine vs volatile anesthetics, rats, 86:141
awareness, with recall, during general anesthesia, incidence and risk factors, 86:1084
cardiac arrest
unsuccessful resuscitation under hypotensive epidural anesthesia during elective hip arthroplasty, 86:847
vasovagal, during insertion of epidural catheter and before administration of epidural medication, 86:1263
cataract formation, posterior subcapsular, associated with multiple lumbar epidural corticosteroid injections, 86:1054
central core disease, patient with, prophylactic use of dantrolene (letter), 86:915
collapsed anterior chamber, during hypothermic CPB (letter), 86:919
complete tracheal disruption, anesthetic management, percutaneous cardiopulmonary support system, 86:998
congenital lobar emphysema, management, use of endobronchial intubation and controlled ventilation, 86:71
delirium, postoperative, development, impact of postoperative pain, 86:781
diabetes mellitus, and difficult laryngoscopy, renal and pancreatic transplant patients, 86:516
diaphragmatic hernia, during pregnancy, management, 86:501
diaphragmatic paralysis, complicating alcohol splanchnic nerve block, 86:845
dural puncture
postdural puncture headache, miscarriage as risk factor (letter), 86:679

prior, effect on CSF sufentanil concentrations after administration of epidural sufentanil, sheep, 86:794
endobronchial intubation, during laparoscopic gynecologic surgery, pneumoperitoneum as risk factor, 86:301
functional impairment, during first seven days after ambulatory surgery, 86:739
gastric insufflation
and LMA (letter), 86:914
risk, and LMA position, 86:867
gastroesophageal reflux, and tracheal contamination, during laparoscopic cholecystectomy and diagnostic gynecological laparoscopy, 86:624
hemodynamic, associated with emergency tracheal reintubation, quality improvement audit, 86:1171
high tracheal stenosis, resection, use of LMA and high-frequency jet ventilation, 86:907
hypercarbia, severe, during CPB, complication of CO_2 flooding surgical field, 86:264
hypertension
acute hemodynamic response during ECT, modification, effectiveness of oral clonidine, 86:1127
response to pin head-holder application during craniotomy, effect of clonidine premedication, 86:1001
hypokalemia, and barbiturate therapy (letter), 86:214
hypotension
anesthesia-induced, patients chronically treated with ACE inhibitors, treatment with angiotensin II, 86:259
cesarean section and bupivacaine baricity, 86:989
propofol-induced, prophylactic effect of ephedrine, elderly female patients, 86:477
refractory, after CPB, treatment with vasopressin, 86:1207
response to pin head-holder application during craniotomy, effect of clonidine premedication, 86:1001
immune thrombocytopenic purpura, parturient with, thromboelastography (letter), 86:675
immunocompromised host, mucormycosis supraglottitis on induction of anesthesia, 86:729
laryngeal carcinoma, undiagnosed, failure of LMA, 86:438
malignant hyperthermia
associated with desflurane anesthesia, humans, 86:1328
suspected, case, during desflurane administration (letter), 86:916
suspected, episode, during desflurane anesthesia (letter), 86:449
syndrome, variant, generalized muscle rigidity and hypercarbia with halothane and isoflurane, 86:297
malignant systemic mastocytosis, patient with, anesthesia using total intravenous technique, 86:442
micrognathia, MRI of airway, infant, 86:964
mucormycosis supraglottitis, on induction of anesthesia, immunocompromised host, 86:729
myxedema coma, treatment, emergency surgery (letter), 86:450
neuroleptic malignant syndrome, severe, efficacy of prolonged large-dose dantrolene, 86:1143
obesity, volume and pH of gastric contents, surgical patients, 86:147
opioid withdrawal syndrome, acute, naloxone-precipitated, after epidural morphine, 86:544
opisthotonos, after flumazenil administered to antagonize midazolam previously administered to treat developing local anesthetic toxicity (letter), 86:677
painful legs and moving toes syndrome, patient with, lumbar sympathetic ganglion block, 86:1056
paresthesia
combined spinal-epidural technique, use of four spinal needles, *in vivo* evaluation, 86:520
during epidural catheter placement, incidence, and combined spinal-epidural analgesia, 86:448
piriformis syndrome, caudal steroids for, mechanism of action (letter), 86:680
pneumothorax, and diaphragmatic hernia, during pregnancy, management, 86:501
pruritus
after administration of 6% hydroxyethyl starch solution, randomized, controlled epidemiology study, 86:850
after cesarean section, intrathecal morphine-induced, subhypnotic doses of propofol do not relieve, 86:310
pseudothrombocytopenia, ethylenediaminetetraacetic acid-dependent, cardiac surgical patient, 86:1210
pulmonary hypertension
chronic hypoxia and monocrotaline injection, effect of prolonged inhaled NO, rats, 86:74
patients with, prostaglandin, route of administration (letter), 86:917
prostaglandin E_1 and right-to-left shunting, 86:703
pyloric stenosis, awake vs paralyzed tracheal intubation, infants, 86:945
retinal hemorrhage, after lumbar epidural steroid injection, 86:119
seizures
acute hemodynamic response during ECT, modification, effectiveness of oral clonidine, 86:1127
lidocaine-induced, dose-dependent suppression by propofol sedation, rats, 86:652
spinal hematoma, spinal and epidural blockade and perioperative low molecular weight heparin, 86:1153
subclinical portal-systemic encephalopathy, patients with, CBF velocity, 86:1005
supraventricular tachydysrhythmias, propofol and, children (letter), 86:914
symptom distress, during first seven days after ambulatory surgery, 86:739
thrombocytopenia, during milrinone infusion after open heart surgery, infants and children, 86:283
torsade de pointes VT and VF, intraoperative, during sevoflurane anesthesia, 86:701
tracheal contamination, and gastroesophageal reflux, during laparoscopic cholecystectomy and diagnostic gynecological laparoscopy, 86:624
transient neurologic syndrome, 1045 patients, after 3% lidocaine spinal anesthesia, 86:1023
trauma, severe, sedating, safety and efficacy of propofol vs midazolam, 86:1219
unplanned extubation, outside OR, quality improvement audit of hemodynamic and tracheal airway complications associated with emergency tracheal reintubation, 86:1171

Complications—*continued*
upper respiratory tract infections, children with, use of LMA, comparison with endotracheal intubation, 86:706
uremic neuropathy, patient with painful legs and moving toes syndrome, lumbar sympathetic ganglion block, 86:1056
von Hippel-Lindau syndrome, excision of cerebellar hemangioblastoma and pheochromocytoma surgery, anesthetic management, 86:673
Compound A, see Metabolism
Computer-assisted auscultation, see Measurement techniques
Confusion assessment method, see Measurement techniques
Congenital lobar emphysema, see Complications
Conjugated estrogen, see Pharmacology
Context-sensitive half-times, see Pharmacokinetics
Continuous spinal anesthesia, see Anesthetic techniques
Coronary artery bypass grafting, see Surgery
Corticosteroids, see Pharmacology
Cost, see Economics
CPB, see Surgery, cardiopulmonary bypass
Craniotomy, see Surgery
Critical care and trauma, see Anesthesia
Cytokines, responses to CPB, effect of CPB duration, 86:28

Dantrolene
see Neuromuscular relaxants
see Pharmacology
Delirium, see Complications
Dental treatment, see Anesthesia
Desflurane, see Anesthetics, volatile
Dexmedetomidine, see Pharmacology
Dextromethorphan, see Pharmacology
Diabetes mellitus, see Complications
Dialyzers, nondialyzability of fentanyl with high-efficiency and high-flux membranes (letter), 86:447
Diaphragm, see Muscle
Diaphragmatic hernia, see Complications
Diaphragmatic paralysis, see Complications
1,2-Dichlorohexafluorocyclobutane, see Nonimmobilizers
2,3-Dichlorooctafluorobutane, see Nonimmobilizers
Diclofenac, see Analgesics
Dimenhydrinate, see Antiemetics
Dobutamine, see Pharmacology
Dopamine, see Pharmacology
Dopexamine, see Pharmacology
Dose-response, see Potency, anesthetic
Double-lumen tube, see Equipment
Droperidol
see Antiemetics
see Pharmacology
Dural puncture, see Complications
Dysrhythmias, see Heart

Economics
and health systems research
aging surgical population and anesthesia workload, 86:1165
costs and new anesthetics, 86:504
diabetes and difficult laryngoscopy, 86:516
economics of clinical pathway for knee surgery, 86:978
emergency tracheal reintubation, -associated complications, quality improvement audit, 86:1171
feedback and cost of anesthesia, 86:510
lidocaine and bowel function, 86:235
multimodal postoperative therapy, 86:228
OR efficiency, 86:896
patient simulators, performance evaluation, testing internal consistency and construct validity, 86:1160
cost
-benefit, effect of timing of ondansetron administration, ambulatory setting, 86:274
comparative, esmolol vs labetalol during ECT (letter), 86:916
-effectiveness, effect of timing of ondansetron administration, ambulatory setting, 86:274
savings, storage period of drug syringes, 86:994
ECT, see Anesthesia, electroconvulsive therapy
Edema, see Brain
Education
patient simulators
competency testing (editorial), 86:1157
performance evaluation, testing internal consistency and construct validity, 86:1160
residents, learning manual skills, recommended number of cases for anesthetic procedures, 86:635
Elastase, see Inflammation
Elastometric balloon pump, see Equipment
Elderly, see Age factors
Electrocardiography, see Monitoring
Electroconvulsive therapy, see Anesthesia
Electroencephalography, see Monitoring
Electrophysiology, action potential and membrane potential, effects of lidocaine, giant axons, crayfish, 86:569
Embolism
CO_2 during hip arthroplasty, infant, 86:299
venous air, continuous, wavelet analysis of embolic heart sound detected by precordial Doppler ultrasound, dogs, 86:325
Emergence, see Recovery
EMLA cream, see Anesthetics, local
Endobronchial intubation
see Complications
see Intubation
Endogenous heparin-like substances, see Blood, coagulation
β-Endorphin, see Pharmacology
Endoscopy mask, see Equipment
Endothelial cells, cultured bovine, production and release of endothelin 1, effect of ketamine, 86:1098
Endotracheal intubation, see Intubation
Endotracheal tube, see Equipment
Enflurane, see Anesthetics, volatile
Enzymes, cholinesterase, plasma cholinesterase activity, decreased, effect on duration of neuromuscular block after mivacurium, postpartum patients, 86:82
Ephedrine, see Sympathetic nervous system, catecholamines
Epidural anesthesia, see Anesthetic techniques
Epinephrine
see Pharmacology
see Sympathetic nervous system, catecholamines
Epithelium, see Airway
Equipment
air release valves, as substitute cufflinks (letter), 86:219
anesthetic machine, absorption of volatile anesthetics, 86:1065
Augustine Guide, blind orotracheal intubation, prospective study, 86:435
catheters
Arrow antecubital catheter, retrieval of air (letter), 86:1337
epidural, continuous, bacterial colonization and infection rate, children, 86:706
epidural, insertion, vasovagal cardiac arrest during, before administration of epidural medication, 86:1263

epidural, intravenous placement, tests to evaluate, prospective clinical study, laboring women, 86:985
epidural, multiport vs uniport (letter), 86:676
Hickman, insertion, radiological screening (letter), 86:216
Combitube, as emergency airway device for anesthesiologists, 86:1141
Datex patient monitoring system, supraventricular tachycardia induced by (letter), 86:1339
double-lumen tube
left-sided, maneuver to position without aid of fiberoptic bronchoscopy, 86:1271
right-sided, vs single-lumen tube with right-sided enclosed bronchial blocker (Univent), for one-lung ventilation, 86:696
tracheal cuff, simple method to protect (letter), 86:675
elastometric balloon pump, postoperative patient-controlled local anesthetic administration, home environment, 86:86
endoscopy mask, jaw support device for fiberscopy, 86:432
endotracheal tube, modified, for infants and small children undergoing video-assisted thoracoscopic surgery, 86:1212
esophageal electrocardiograph, detection of myocardial ischemia, 86:34
fiberoptic bronchoscope, jaw support device for fiberscopy, 86:432
HealthQuiz, vs written questionnaire, history-taking process during preanesthetic visit, 86:134
jaw support device, facilitation of laryngeal exposure and ventilation during fiberoptic intubation, 86:432
laryngeal mask airway
and high-frequency jet ventilation, for resection of high tracheal stenosis, 86:907
failure, undiagnosed laryngeal carcinoma, 86:438
gastric insufflation and (letter), 86:914
insertion techniques (letter, reply), 86:1337, 1338
position, and risk of gastric insufflation, 86:867
use in children with upper respiratory tract infections, comparison with endotracheal intubation, 86:706
vs cuffed tracheal tube, bronchial mucus velocity in paralyzed anesthetized patients, 86:1280
laryngoscope blade, failure (letter), 86:448
Luer slip connectors, as substitute, studs (letter), 86:219
needles
gauge, effect on pain during intradermal injection, 86:379
pencil-point spinal, used for combined spinal-epidural technique, *in vivo* evaluation, 86:520
single-lumen tube, with right-sided enclosed bronchial blocker (Univent), vs right-sided double-lumen tube, for one-lung ventilation, 86:696
stethoscope, esophageal, placement depth, effect on heart and lung sound monitoring during general anesthesia, 86:1276
syringes, anesthetic and resuscitative drug, used in obstetric OR, sterility, 86:994
TEE probe, simple holder (letter), 86:677
tracheal tube, cuffed, vs LMA, bronchial mucus velocity in paralyzed anesthetized patients, 86:1280
Esmolol, see Pharmacology
Esophageal electrocardiograph, see Equipment
Esophagectomy, see Surgery
Ethics, human toxicity study (letter), 86:445
Ethylenediaminetetraacetic acid, see Pharmacology
Etomidate, see Anesthetics, intravenous
Evaluation, performance, patient simulator, testing internal consistency and construct validity, 86:1160
Extracorporeal shock wave lithotripsy, see Surgery
Eye
anterior chamber, collapsed, during hypothermic CPB (letter), 86:919
eyelid movement, during complete neuromuscular block, 86:1332
intraocular pressure, effect of pneumoperitoneum, α-chymotrypsin-induced glaucoma, rabbits, 86:1283
retinal hemorrhage, after lumbar epidural steroid injection, 86:119

Fascia iliaca compartment block, see Anesthetic techniques
Fast computed tomography, see Measurement techniques
Fast-track eligibility, see Recovery
Femoral three-in-one nerve block, see Anesthetic techniques, three-in-one block
Fentanyl
see Analgesics
see Anesthetics, intravenous
see Pharmacokinetics
Fentanyl Oralet, see Premedication, oral transmucosal fentanyl citrate
Fiberoptic bronchoscope, see Equipment
Fiberscopy, see Anesthetic techniques
FK409, see Gases, nonanesthetic, nitric oxide
Flow-volume curves, see Monitoring
Fluids, intravenous, effects on brain edema two hours after closed head trauma, rats, 86:1225
Flumazenil, see Pharmacology
Fluoride, see Ions
Formalin test, see Pain
Functional impairment, see Complications

Gabapentin, see Analgesics
Gases, nonanesthetic, nitric oxide
effects on IL-8 and elastase from lipopolysaccharide-stimulated whole blood, 86:427
FK409, and peripherally administered morphine, interaction, during experimental inflammation, 86:367
inhaled, prolonged, attenuation of hypoxic pulmonary vascular remodeling, rats, 86:74
intrinsic, effects on cardiac neural regulation, cats, 86:1194
peroxynitrate, effects on IL-8 and elastase from lipopolysaccharide-stimulated whole blood, 86:427
placental tissue enhancement of uterine relaxation by nitroglycerin, 86:304
synthase
alveolar macrophages, effect of dantrolene, rats, 86:1065
neuronal, inhibitor, 7-nitroindazole, selectivity (letter), 86:679
Gastric contents, volume and pH, obese vs lean surgical patients, 86:147
Gastric insufflation, see Complications
Gastric mucosal oxygen delivery, during CPB, decrease, despite constant systemic oxygen delivery, 86:455
Gastroesophageal reflux, see Complications
Gastrointestinal tract
bowel function, faster return, lidocaine-treated patients undergoing radical retropubic prostatectomy, 86:235
motility, epidural anesthesia and, 86:837

General anesthesia, see Anesthetic techniques
Gender
differences, nociception and response to ibuprofen, 86:1257
effects on optimal premedication dose of intramuscular midazolam, 86:1103
pharmacokinetics of vecuronium, male and female patients, 86:1322
Gentamicin, see Antibiotics
Glibenclamide, see Pharmacology
Glossopharyngeal nerve block, see Anesthetic techniques
Glucagon, see Pharmacology
Glutamate, see Amino acids, excitatory
Granisetron, see Antiemetics
Gynecological surgery, see Surgery

Halothane, see Anesthetics, volatile
Hand surgery, see Surgery
HealthQuiz, see Equipment
Heart
aortic stenosis, patients with, intrathecal sufentanil analgesia for extracorporeal shock wave lithotripsy, 86:943
arrhythmias
abnormal automaticity, effects of halothane, cardiac Purkinje fibers, dogs, 86:488
bupivacaine-induced, effects of verapamil and nimodipine, *in vivo* and *in vitro* study, rats, 86:818
torsade de pointes VT and VF, intraoperative, during sevoflurane anesthesia, 86:701
atrial septal defect, and predominant right-to-left shunting, preoperative prostaglandin E_1 therapy, 86:703
atrial trabeculae, contractile force, effect of oxytocin and its preservative, 86:40
dysrhythmias
perioperative ventricular, patients with heart disease undergoing noncardiac surgery, 86:16
supraventricular tachydysrhythmias, propofol and, children (letter), 86:914
ischemia
detection, use of new high-resolution esophageal electrocardiography recording technique, 86:34
reperfusion, cardioprotective effects of propofol, rats, 86:252
left ventricle
afterload, determinants, adverse alteration by etomidate, dogs with dilated cardiomyopathy, 86:932
loading modifications, induced by pneumoperitoneum, time course echocardiographic study, 86:482
systolic and diastolic function, unaltered during propofol infusion, newborn swine, 86:717
myocardial depression, effects of intravenous levobupivacaine and bupivacaine, sheep, 86:797
neural regulation, effects of intrinsic NO, cats, 86:1194
output
and onset time of different muscle relaxants (letter), 86:1146
effect of sevoflurane, dogs, 86:1079
Purkinje fibers, effects of halothane on abnormal automaticity, dogs, 86:488
rate
effect of sevoflurane, dogs, 86:1079
effects of cisatracurium vs atracurium, neurosurgical patients, 86:123
influence of bolus injection rate of propofol, sheep, 86:1109
reversed shunting, and atrial septal defect, patient with, preoperative prostaglandin E_1 therapy, 86:703
tachycardia, supraventricular, induced by Datex patient monitoring system, 86:1339
ventricular dysfunction, amrinone vs milrinone after cardiac surgery, 86:683
Hemidiaphragmatic paresis, see Ventilation
Hemodilution, see Blood
Hemodynamics
acute hemodynamic response during ECT, modification, effectiveness of oral clonidine, 86:1127
changes, sevoflurane-associated, dogs, 86:1079
effects of anesthesia, patients receiving angiotensin-converting enzyme inhibitors (letter), 86:215
gasless mechanical vs conventional CO_2 pneumoperitoneum methods for laparoscopic cholecystectomy, 86:153
Heparin
see Blood, coagulation, endogenous heparin-like substances
see Pharmacology
Hernia repair, see Surgery
High tracheal stenosis, see Complications
Hip surgery, see Surgery
History-taking methods, see Preoperative evaluation
Hormones
oxytocin, effect on contractile force of human atrial trabeculae, 86:40
stress, neuroendocrine, failure to recreate postoperative hypercoagulable state, 86:640
thyroid, modulation, effects on liver injury associated with ischemia-reperfusion and cold storage, rats, 86:1187
Hospitals
anesthesia workload, influence of aging surgical population, ten-year survey, 86:1165
costs, perioperative clinical pathway for knee replacement surgery, clinical implications for ambulatory surgery, 86:978
length-of-stay, decreased, lidocaine-treated patients undergoing radical retropubic prostatectomy, 86:235
Hydroxyethyl starch
see Blood, plasma substitutes
see Pharmacokinetics
Hyperalgesia, see Pain
Hypercapnia, see Carbon dioxide
Hypercarbia
see Carbon dioxide
see Complications
Hypertension, see Complications
Hyperthermia, see Complications, malignant hyperthermia
Hypnotics, chloral hydrate, sedation, additive sedative and respiratory depressant effects of N_2O, 86:724
Hypotension
see Blood pressure
see Complications
Hypothermia, see Temperature
Hypothermic CPB, see Surgery, cardiopulmonary bypass
Hypoxia
acute and sustained, ventilatory responses, during isoflurane anesthesia, 86:403
chronic, pulmonary vascular remodeling, attenuation, effectiveness of prolonged inhaled NO, rats, 86:74

Ibuprofen, see Analgesics
Idazoxan, see Anesthetics, intravenous
Immune response
immunoglobins, influence of prolonged oral treatment with sustained-release morphine tablets, 86:166
immunosuppression, effect of midazolam and propofol on IL-8

from human polymorphonuclear leukocytes, 86:1289
lymphocytes
influence of prolonged oral treatment with sustained-release morphine tablets, 86:166
peripheral blood B lymphocytes, chemotaxis by β-endorphin, reversibility by naloxone, 86:670
Immune thrombocytopenic purpura, see Complications
Immunocompromised host, see Complications
Infection
bacterial growth, *Pseudomonas aeruginosa,* inhibition by halogenated anesthetics, 86:1075
continuous epidural catheters, children, 86:712
sterility
of drugs in syringes for long times, 86:994
of thiopental after preparation, 86:208
Inflammation
elastase, from lipopolysaccharide-stimulated whole blood, effects of NO and peroxynitrate, 86:427
experimental, interaction of FK409 and peripherally administered morphine, 86:367
IL-6, response to cardiac surgery, influence of duration of CPB or aortic cross-clamp in absence of blood and/or blood product administration, 86:28
IL-8, from lipopolysaccharide-stimulated whole blood, effects of NO and peroxynitrate, 86:427
systemic inflammatory response to CPB, manifestations, 86:22
Informed consent, see Research
Injection pain, see Pain
Inositol-triphosphate, see Analgesics
Interactions, drug
alfentanil, and ketamine, 86:1250
atracurium, and mivacurium, 86:62
cisatracurium, and mivacurium, 86:62
clonidine, and epinephrine, 86:1010
epinephrine, and clonidine, 86:1010
fentanyl, and propofol, 86:872
FK409, and morphine, 86:367
ketamine, and alfentanil, 86:1250
lidocaine, and propofol, 86:652
mivacurium
and atracurium, 86:62
and cisatracurium, 86:62
morphine, and FK409, 86:367
propofol
and fentanyl, 86:872
and lidocaine, 86:652
and thiopental, 86:422
thiopental, and propofol, 86:422
Interleukin-6, see Inflammation
Interleukin-8
see Inflammation
see Pharmacology
Interscalene block, see Anesthetic techniques
Intravenous regional anesthesia, see Anesthetic techniques
Intubation
difficult, renal and pancreatic transplant patients with diabetes mellitus, 86:516
endobronchial, and controlled ventilation, management of congenital lobar emphysema, 86:71
endotracheal, with laryngoscope or fiberscope, required propofol concentration, 86:872
orotracheal, blind, with Augustine Guide, prospective study, 86:435
tracheal
ambulatory surgery patients, using remifentanil and propofol without muscle relaxants, 86:45
circulatory responses, blunting, optimal injection time of small-dose fentanyl, 86:658
comparison with LMA, children with upper respiratory tract infections, 86:706
LMA position and risk of gastric insufflation, 86:867
reintubation, emergency, -associated hemodynamic and tracheal airway complications, quality improvement audit, 86:1171
technique, awake vs paralyzed, infants with pyloric stenosis, 86:945
technique, emergency, use of Combitube, 86:1137
technique, fiberoptic, facilitation of laryngeal exposure and ventilation, use of jaw support device, 86:432
technique, rocuronium using "timing principle," 86:1137
Ions
calcium, channels, single human neuronal L-type, effects of halothane, 86:885
fluoride, prolonged administration of isoflurane, patients with severe renal dysfunction, 86:440
magnesium
-treated parturients, mivacurium infusion rate and spontaneous neuromuscular recovery, 86:523
vs fentanyl, anesthetic supplementation, sensory changes and pain after abdominal hysterectomy, 86:95
Ischemia
see Brain
see Heart
see Liver
Ischemic insult, see Kidney
Isoflurane, see Anesthetics, volatile
Isoproterenol, see Pharmacology

Jaw support device, see Equipment
Jet ventilation, see Ventilation

Ketamine, see Anesthetics, intravenous
Ketorolac, see Analgesics
Kidney
dysfunction
perioperative, ketorolac-associated, factors influencing, 86:1090
preoperative, patients undergoing CPB, renal protection, 86:3
severe, patients with, prolonged administration of isoflurane, 86:440
function, gasless mechanical vs conventional CO_2 pneumoperitoneum methods for laparoscopic cholecystectomy, 86:153
ischemic insult, kidney viability, and renal function (editorial), 86:1
toxicity, absence, after four hours of 1.25 MAC sevoflurane anesthesia, volunteers, 86:662
Knee surgery, see Surgery

Labetalol, see Pharmacology
Lactate, see Metabolism
Laminectomy, see Surgery
Laparoscopic surgery
see Surgery, cholecystectomy
see Surgery, orchidopexy
see Surgery, gynecological
Laparotomy, see Surgery
Laryngeal carcinoma, see Complications
Laryngeal mask airway, see Equipment
Laryngectomy, see Surgery
Laryngoscope blade, see Equipment
Laryngoscopy, see Anesthetic techniques
Latex, see Allergy
Length-of-stay, see Hospitals
Levobupivacaine, see Anesthetics, local
Lidocaine
see Anesthetics, intravenous
see Anesthetics, local
Lipid peroxidation, see Metabolism
Lipids, see Brain

Lipopolysaccharide, -stimulated whole blood, IL-8 and elastase, effects of NO and peroxynitrate, 86:427
Liver
blood flow, influences of CPB and fentanyl anesthesia, beagles, 86:1177
disease, chronic, CBF velocity, 86:1005
injury, associated with ischemia-reperfusion and cold storage, effects of thyroid hormone modulation, rats, 86:1187
ischemia, -reperfusion and cold storage, injury associated with, effects of thyroid hormone modulation, rats, 86:1187
oxygen metabolism, influences of CPB and fentanyl anesthesia, beagles, 86:1177
toxicity, absence, after four hours of 1.25 MAC sevoflurane anesthesia, volunteers, 86:662
Liver transplantation, see Surgery
LMA, see Equipment, laryngeal mask airway
Lobectomy, see Surgery
Lornoxicam, see Analgesics
Luer slip connectors, see Equipment
Lumbar disk surgery, see Surgery
Lumbar sympathetic ganglion block, see Anesthetics techniques, sympathetic block
Lumbosacral dermatomes, duration of lidocaine spinal anesthesia, effect of varied doses of epinephrine, 86:1018
Lung
alveolar macrophages, L-arginine transport and NO synthase, effect of dantrolene, rats, 86:1065
aspiration, gastric contents and obesity, 86:141
bronchial mucus transport velocity, paralyzed anesthetized patients, LMA vs cuffed tracheal tube, 86:1280
bronchodilation, effect of volatile anesthetics, epithelial dependence, distal bronchi, rats, 86:646
chronic obstructive pulmonary disease
minimally invasive myocardial revascularization before lung transplantation, anesthetic management, 86:939
successful continuous spinal anesthetic, 86:591
dynamic hyperinflation, intrinsic PEEP and flow-volume curves, 86:880
functional residual capacity, pulmonary vagal afferents and respiratory distress, halothane-anesthetized cats, 86:398
injury
acid-induced, after general anesthesia, effects of volatile anesthetics vs ketamine, rats, 86:141
acute, in severe multiple organ failure, treatment with inhaled prostaglandin E_1, 86:753
mechanics, diaphragmatic function, effects of cervical epidural anesthesia, conscious patients, 86:1033
pulmonary arterial concentrations of bupivacaine enantiomers after epidural injection, elderly patients, 86:812
pulmonary gas exchange, effect of dopexamine, anesthetized, paralyzed patients, 86:314
Lung transplantation, see Surgery
Lymphocytes, see Immune response

MAC, see Potency, anesthetic
Magnesium, see Ions
Magnesium sulfate, see Receptors, NMDA, antagonists
Magnetic resonance imaging, see Measurement techniques
Malignant hyperthermia, see Complications
Malignant systemic mastocytosis, see Complications
Mannitol, see Pharmacology
Measurement techniques
capnography, isocapnic hypoxic ventilatory responses during isoflurane anesthesia, 86:403
computer-assisted auscultation, esophageal stethoscope placement depth, effect on heart and lung sound monitoring during general anesthesia, 86:1276
confusion assessment method, postoperative pain and delirium, 86:781
fast computed tomography, effect of N_2O on chest wall function, humans and dogs, 86:1058
MAC for airway occlusion, halothane-anesthetized cats, 86:398
multiple inert gas elimination technique, determination of ventilation perfusion distribution and pulmonary gas exchange, effects of dopexamine, anesthetized, paralyzed patients, 86:314
laser Doppler flowmetry
gastric mucosal red blood cell flux during CPB, 86:455
skin vasomotor reflex, assessment of level of regional anesthesia, 86:336
MRI, airway, infant with micrognathia, 86:964
pneumotachography, isocapnic hypoxic ventilatory responses during isoflurane anesthesia, 86:403
power spectrum analysis, etomidate and aortic input impedance, 86:932
pulse oximetry, signal improvement, use of EMLA cream (letter), 86:915
respiratory impedance plethysmography, effect of N_2O on chest wall function, humans and dogs, 86:1058
respiratory inductive plethysmography, assessment of respiratory function after supraclavicular block, 86:1239
visual analog scale
immediate postoperative period, intrasubject variability and correlation with numeric scale, 86:102
postoperative pain and delirium, 86:781
Mechanisms of anesthetic action, ventilatory effects of nonimmobilizers, swine, 86:173
Meperidine, see Analgesics
Mepivacaine, see Anesthetics, local
Meta-analysis, see Statistics
Metabolism
compound A, kinetic characteristics, effect of anesthetic duration, volunteers, 86:414
lactate, during normothermic CPB, humans, 86:22
lipid peroxidation, during reperfusion, attenuation by propofol, ischemic reperfused isolated rat hearts, 86:252
Micrognathia, see Complications
Midazolam, see Anesthetics, intravenous
Milrinone
see Pharmacokinetics
see Pharmacology
Minimum alveolar concentration, see Potency, anesthetic
Miscarriage, as risk factor for postdural puncture headache (letter), 86:679
Mivacurium, see Neuromuscular relaxants
Mivazerol, see Anesthetics, intravenous
MK801, see Receptors, NMDA, antagonists
Monitored anesthesia care
changing role, ambulatory setting (letter, replies), 86:1335, 1336

dissociative, not cited (letter, reply), 86:1336
Monitoring
aortic blood flow echo-Doppler, continuous esophageal, assessment of hemodynamic effects of pneumoperitoneum during laparoscopic surgery, healthy infants, 86:283
CO_2
noninvasive techniques, end-tidal vs transcutaneous CO_2 (letter), 86:675
tensions, during anesthesia in prone position, 86:668
central venous pressure, effect of pneumoperitoneum on intraocular pressure, α-chymotrypsin-induced glaucoma, rabbits, 86:1283
EEG
burst suppression during CPB, thiopental vs isoflurane, 86:246
quantitative topographical analysis after intravenous clonidine, healthy volunteers, 86:202
electrocardiography, esophageal, new high-resolution recording technique, detection of myocardial ischemia, 86:34
flow-volume curves, and intrinsic PEEP, 86:880
precordial Doppler ultrasound, embolic heart sound during continuous venous air embolism, wavelet analysis, dogs, 86:325
thromboelastography
heparin activity during liver transplantation, 86:691
parturient with immune thrombocytopenic purpura (letter), 86:675
transcranial Doppler ultrasonography
CBF velocity and cerebral oxygen extraction during CPB, effects of propofol, 86:1201
CBF velocity in chronic liver disease, 86:1005
effect of hemodilution on CBF velocity, anesthetized patients, 86:320
effects of cisatracurium vs atracurium, neurosurgical patients, 86:123
transesophageal echocardiography (TEE)
probe, simple holder (letter), 86:677
time course study, left ventricular loading modifications induced by pneumoperitoneum, 86:482
treating anesthesia-induced hypotension with angiotensin II, patients chronically treated with ACE inhibitors, 86:259
Morphine, see Analgesics
MRI, see Measurement techniques, magnetic resonance imaging
Mucormycosis supraglottitis, see Complications
Multiple inert gas elimination technique, see Measurement techniques
Muscle
chest wall, function, effect of N_2O, humans and dogs, 86:1058
diaphragm, contractility, fatigue-induced changes, effect of halothane and sevoflurane, hamsters, 86:392
rigidity, generalized, and hypercarbia, with halothane and isoflurane, 86:297
sternocleidomastoid, as surface landmark for supraclavicular nerve blocks (letter), 86:1149
Myxedema coma, see Complications

Naloxone, see Antagonists, opioid
Needles, see Equipment
Neostigmine, see Analgesics
Nerve
cardiac sympathetic nerve activity, effects of intrinsic NO, cats, 86:1194
injury, -induced pain, effects of dexmedetomidine, spinal nerve ligation model, rats, 86:355
stimulation, nonadrenergic and noncholinergic, relaxant response of isolated cerebral arteries, effects of halothane vs isoflurane, dogs, 86:552
Neuroleptic malignant syndrome, see Complications
Neuromuscular relaxants
atracurium
pharmacodynamics of mivacurium preceded by, children, 86:62
vs cisatracurium, cerebral and cardiovascular effects, neurosurgical patients, 86:123
choice, and postoperative recovery time (letter, reply), 86:1335
cisatracurium
pharmacodynamics of mivacurium preceded by, children, 86:62
vs atracurium, cerebral and cardiovascular effects, neurosurgical patients, 86:123
dantrolene, prolonged large-dose, efficacy for severe neuroleptic malignant syndrome, 86:1143
mivacurium
duration of neuromuscular block, effect of decreased plasma cholinesterase activity, postpartum patients, 86:82
infusion rate and spontaneous neuromuscular recovery, magnesium-treated parturients, 86:523
preceded by atracurium or cisatracurium, pharmacodynamics, children, 86:62
onset time, and cardiac output (letter), 86:1146
rocuronium, tracheal intubation using "timing principle," 86:1137
succinylcholine
and desflurane, malignant hyperthermia, humans, 86:1328
tracheal intubation, vs tracheal intubation with rocuronium using "timing principle," 86:1137
vecuronium
dose-response and time course of action, influence of acute normovolemic hemodilution, 86:861
eyelid movement during complete neuromuscular block, 86:1332
pharmacokinetics, male and female patients, 86:1322
Neurons, single human neuronal L-type calcium channels, effects of halothane, 86:885
Neurosurgical anesthesia, see Anesthesia
Neurotoxicity, see Toxicity
Neutrophil functions, human, inhibitory effects of intravenous anesthetics, 86:159
Nifedipine, see Pharmacology
Nimodipine, see Pharmacology
Nitric oxide, see Gases, nonanesthetic
Nitroglycerin, see Pharmacology
7-Nitroindazole, see Pharmacology
Nitrous oxide, see Anesthetics, gases
NMDA, see Receptors
Noncardiac surgery
see Outcome
see Surgery
Nonimmobilizers
1,2-dichlorohexafluorocyclobutane, ventilatory effects, swine, 86:173
2,3-dichlorooctafluorobutane, ventilatory effects, swine, 86:173
lack of effect on nociceptive responses, 86:629
perfluoropentane, ventilatory effects, swine, 86:173
Nonsteroidal antiinflammatory drugs, see Analgesics

Nose, alae nasi, inspiratory collapse, cause of variable extrathoracic airway obstruction, 86:294
NSAIDs, see Analgesics, nonsteroidal antiinflammatory drugs

Obesity, see Complications
Observational studies, vs randomized controlled trials (editorial), 86:225
Obstetric anesthesia, see Anesthesia
Oleamide, see Brain, lipids
Ondansetron, see Antiemetics
One-lung ventilation, see Anesthetic techniques
Open heart surgery, see Surgery
Operating room (OR)
efficiency, improving, successful strategies, academic institutions, 86:896
obstetric, sterility of anesthetic and resuscitative drug syringes, 86:994
Opioid withdrawal syndrome, see Complications
Opisthotonos, see Complications
Oral transmucosal fentanyl citrate, see Premedication
Orchidopexy, see Surgery
Orotracheal intubation, see Intubation
Outcome
noncardiac surgery, patients with heart disease, incidence of perioperative dysrhythmias, 86:16
pulmonary, comparative effects of postoperative analgesic therapies, cumulative meta-analyses of randomized, controlled trials, 86:598
Oxygen
consumption, changes after tourniquet release, patients breathing spontaneously under epidural anesthesia, 86:90
delivery, systemic, during CPB, decreased gastric mucosal oxygen delivery, 86:455
extraction, cerebral, during CPB, effects of propofol, 86:1201
metabolism, hepatic, influences of CPB and fentanyl anesthesia, beagles, 86:1177
transport, splanchnic, during normothermic CPB, humans, 86:22
Oxygenation
arterial, one-lung ventilation, isoflurane vs sevoflurane, 86:1266
regional cerebral, differential effects of prostaglandin E_1 and nitroglycerin (letter, reply), 86:1338, 1339
Oxytocin
see Hormones
see Pharmacology

Pain
antinociception
effect of desflurane, 86:629
effect of morphine, potentiation, intrathecal co-infusion of magnesium sulfate, rats, 86:830
effect of morphine, potentiation, ketamine vs dextromethorphan, rats, 86:825
effect of N_2O, 86:629
effect of S(-)-enantiomer of ketamine on carrageenan hyperalgesia after intrathecal administration, rats, 86:561
cancer, epidural NSAIDs, 86:117
chronic
intractable, treatment with topical large-dose capsaicin, preliminary report, 86:579
prolonged oral treatment with sustained-release morphine tablets, influence on immune function, 86:166
control, home environment, postoperative patient-controlled local anesthetic administration, 86:86
formalin test, experimental inflammation, interaction of FK409 and peripherally administered morphine, 86:367
hyperalgesia
after thermal injury, effect of intrathecal gabapentin and 3-isobutyl γ-aminobutyric acid, rats, 86:348
effect of desflurane, 86:629
pinprick, effects of intravenous ketamine-alfentanil combinations vs either drug alone, normal subjects, 86:1250
injection
lidocaine, effect of needle gauge and lidocaine pH, 86:379
propofol, opioid peripheral analgesic effects, 86:382
neuropathic
allodynia, mechanical, effect of ECT, rat model of peripheral neuropathy, 86:584
allodynia, produced by administration of anti-GD_2 ganglioside, reversal, effectiveness of gabapentin, 86:111
hyperalgesia, mechanical, delaying effect of preemptive intrathecal ketamine, rats, 86:557
hyperalgesia, thermal, effect of ECT, rat model of peripheral neuropathy, 86:584
spinal nerve ligation model, effects of dexmedetomidine, rats, 86:355
peripheral, mediation of analgesia by neostigmine, inflamed knee joint, rats, 86:1027
postoperative, management
alfentanil as adjuvant to epidural bupivacaine, lack of evidence of spinal action, laparotomy, 86:574
analgesic therapies, comparative effects on pulmonary outcome, meta-analyses of randomized, controlled trials, 86:598
comparison of anesthetic supplementation with fentanyl vs magnesium or ketamine, abdominal hysterectomy, 86:95
continuous pediatric epidural catheters, 86:712
efficacy, assessment, visual analog scale, intrasubject variability and correlation with numeric scale, 86:102
epidural verapamil, lower abdominal surgery, 86:786
impact on development of postoperative delirium, 86:781
intravenous lidocaine, patients undergoing radical retropubic prostatectomy, 86:235
multimodal, abdominothoracic esophagectomy, 86:228
multimodal, anterior cruciate ligament repair, 86:374
multimodal, PCEA regimen, effect of adding ketamine, 86:1245
premedication with oral dextromethorphan, tonsillectomy, 86:594
small doses of intrathecal morphine combined with systemic diclofenac, cesarean section, 86:538
thermal injury, hyperalgesia after, effect of intrathecal gabapentin and 3-isobutyl γ-aminobutyric acid, rats, 86:348
tolerance
effects of intravenous ketamine-alfentanil combinations vs either drug alone, normal subjects, 86:1250
gender differences, nociception and response to ibuprofen, 86:1257
tourniquet, epinephrine and bupivacaine spinal anesthesia, 86:973
vascular, acute, failure of sympathetic block to reduce, humans, 86:588
Painful legs and moving toes syndrome, see Complications
Paresthesia, see Complications

Patient-controlled analgesia, see Anesthetic techniques
Patient-controlled epidural analgesia, see Anesthetic techniques
Patient simulators, see Education
PCA, see Anesthetic techniques, patient-controlled analgesia
PCEA, see Anesthetic techniques, patient-controlled epidural analgesia
Pediatric anesthesia, see Anesthesia
Pentobarbital, see Anesthetics, intravenous
Percutaneous cardiopulmonary support system, see Anesthetic techniques
Perfluoropentane, see Nonimmobilizers
Pharmacodynamics
context-sensitive half-times, clinical implication, recovery from opioid anesthesia, 86:184
mivacurium
metabolism by plasma cholinesterase, postpartum patients, 86:82
preceded by atracurium or cisatracurium, children, 86:62
propofol, injection rate, influence on cardiovascular effects and peak blood concentrations, sheep, 86:1109
vecuronium, influence of acute normovolemic hemodilution, 86:861
Pharmacokinetics
acetyl starch, as plasma volume expander, patients undergoing elective surgery, 86:856
alfentanil, as adjuvant to epidural bupivacaine, management of postoperative pain after laparotomy, lack of evidence of spinal action, 86:574
bupivacaine enantiomers
after epidural administration of racemates, 86:361
systemic and regional, sheep, 86:805
clearance, morphine, postoperative infants during intravenous infusion, influence of age and surgery, 86:958
compound A, effect of anesthetic duration, volunteers, 86:414
context-sensitive half-times, clinical implication, recovery from opioid anesthesia, 86:184
desflurane, vs sevoflurane, effect of anesthetic duration, volunteers, 86:414
fentanyl, nondialyzability with high-efficiency and high-flux membranes (letter), 86:447
hydroxyethyl starch, as plasma volume expander, patients undergoing elective surgery, 86:856
mepivacaine enantiomers, after epidural administration of racemates, 86:361
milrinone, after open heart surgery, infants and children, 86:283
ocular transmucosal, absorption and toxicity of sufentanil, dogs, 86:138
oral mucosa, buccal absorption of etomidate from solid formulation, dogs, 86:1116
oral transmucosal fentanyl citrate, premedication with, pediatric tonsillectomy patients, 86:66
propofol, rate of administration, effect on brain concentrations, sheep, 86:1302
sevoflurane, vs desflurane, effect of anesthetic duration, volunteers, 86:414
vecuronium, male and female patients, 86:1322
Pharmacology
adenosine triphosphate, effect on sevoflurane requirements for MAC and MAC-awake, 86:179
amrinone
use after cardiac valve surgery, 86:461
vs milrinone, after elective cardiac surgery, multicenter, randomized, blind comparison, 86:683
ACE inhibitors
patients chronically treated with, anesthesia-induced hypotension, treatment with angiotensin II, 86:259
patients receiving, hemodynamic effects of anesthesia (letter), 86:215
angiotensin II
for treatment of anesthesia-induced hypotension, patients chronically treated with ACE inhibitors, 86:259
infusion of stress hormones, failure to recreate postoperative hypercoagulable state, 86:640
clonidine
epidural, spinal cord accumulation, effects of epidural epinephrine, rabbits (letter), 86:1145
intravenous, quantitative topographical EEG analysis after, healthy male volunteers, 86:202
oral, effectiveness in modifying acute hemodynamic response during ECT, 86:1127
oral, premedication, reduction of awakening concentration of isoflurane, 86:410
topical and intravenous, vascular effects, canine pial microcirculation, 86:766
conjugated estrogen, effects on transfusion and coagulation factor requirements, orthotopic liver transplantation, 86:1183
corticosteroids
cortisol, infusion of stress hormones, failure to recreate postoperative hypercoagulable state, 86:640
injections, multiple lumbar epidural, posterior subcapsular cataract formation associated with, 86:1054
dantrolene
inhibition of L-arginine transport and NO synthase, alveolar macrophages, rats, 86:1065
prophylactic use, patient with central core disease (letter), 86:915
dexmedetomidine, effects in spinal nerve ligation model of neuropathic pain, rats, 86:355
dextromethorphan
oral, premedication with, reduction of postoperative pain after tonsillectomy, 86:594
vs ketamine, potentiation of antinociceptive effect of morphine, rats, 86:825
dobutamine, use after cardiac valve surgery, 86:461
dopamine
renal protection in cardiac surgery, 86:3
use after cardiac valve surgery, 86:461
dopexamine, effect on ventilation perfusion distribution and pulmonary gas exchange, anesthetized, paralyzed patients, 86:314
droperidol
and side effects of epidural morphine after cesarean section, 86:532
vs ondansetron, efficacy, safety, and patient satisfaction as antiemetics for elective outpatient surgical procedures, 86:731
β-endorphin, chemotaxis of human peripheral blood B lymphocytes, reversibility by naloxone, 86:670
epinephrine
and isoproterenol, combination, as simulated epidural test dose, isoflurane-anesthetized adults, 86:1312
effect on small-dose hyperbaric bupivacaine spinal anesthesia, clinical implications for ambulatory surgery, 86:973

Pharmacology—*continued*
epidural, effects on spinal cord accumulation of epidural clonidine, rabbits (letter), 86:1145
infusion of stress hormones, failure to recreate postoperative hypercoagulable state, 86:640
0.125%, addition of isoproterenol, effects on quality and duration of epidural analgesia, laboring parturients, 86:749
use after cardiac valve surgery, 86:461
varied doses, effect on duration of lidocaine spinal anesthesia, thoracic and lumbosacral dermatomes, 86:1018
esmolol, vs labetalol, during ECT, comparative doses and cost (letter), 86:916
ethylenediaminetetraacetic acid, -dependent thrombocytopenia, cardiac surgical patient, 86:1210
flumazenil, administered to antagonize midazolam previously administered to treat developing local toxicity, triggering of opisthotonos (letter), 86:677
glibenclamide, clonidine effects on pial microcirculation, 86:766
glucagon, infusion of stress hormones, failure to recreate postoperative hypercoagulable state, 86:640
heparin, low molecular weight, perioperative, spinal and epidural blockade (editorial), 86:1153
IL-8, from human polymorphonuclear leukocytes, effect of midazolam and propofol, 86:1289
isoproterenol
addition to 0.125% bupivacaine, effects on quality and duration of epidural analgesia, laboring parturients, 86:749
epinephrine and, combination, as simulated epidural test dose, isoflurane-anesthetized adults, 86:1312
labetalol, vs esmolol, during ECT, comparative doses and cost (letter), 86:916
mannitol, hyperosmolar, intracarotid injection, disruption of blood-brain barrier, effects of pentobarbital, rats, 86:1230
midazolam, administered to treat developing local anesthetic toxicity, flumazenil administered to antagonize, triggering of opisthotonos (letter), 86:677
milrinone, vs amrinone, after elective cardiac surgery, multicenter, randomized, blind comparison, 86:683
nifedipine, effect of ECT on thermal hyperalgesia and mechanical allodynia, rat model of peripheral neuropathy, 86:584
nimodipine, effects on bupivacaine-induced cardiotoxicity, *in vivo* and *in vitro* study, rats, 86:818
nitroglycerin
and prostaglandin E_1, differential effects on regional cerebral oxygenation (letter, reply), 86:1338, 1339
-induced vasodilation of pial arteries, effect of mild hypothermia, cats, 86:546
uterine relaxation, enhancement by placental tissue, 86:304
7-nitroindazole, selectivity (letter), 86:679
ondansetron, vs droperidol, efficacy, safety, and patient satisfaction as antiemetics for elective outpatient surgical procedures, 86:731
oxytocin, intravenous, patients undergoing elective cesarean section (letter, reply), 86:1334
prostaglandin E_1
and nitroglycerin, differential effects on regional cerebral oxygenation (letter, reply), 86:1338, 1339
inhaled, treatment of acute lung injury in severe multiple organ failure, 86:753
therapy, patient with atrial septal defect and predominant right-to-left shunting, 86:703
riluzole, failure to attenuate increases in hippocampal glutamate concentrations during repeated transient global cerebral ischemia, rabbits, 86:128
steroids, caudal, for piriformis syndrome, mechanism of action (letter), 86:680
vasopressin
infusion of stress hormones, failure to recreate postoperative hypercoagulable state, 86:640
treatment of refractory hypotension after CPB, 86:1207
verapamil
effects on bupivacaine-induced cardiotoxicity, *in vivo* and *in vitro* study, rats, 86:818
epidural, reduction of analgesic consumption after lower abdominal surgery, 86:786
yohimbine, clonidine effects on pial microcirculation, 86:766
Pheochromocytoma surgery, see Surgery
Pial arteries, see Arteries
Piriformis syndrome, see Complications
Plasma substitutes, see Blood
Platelets, see Blood
Pneumoperitoneum, see Carbon dioxide, insufflation
Pneumotachography, see Measurement techniques
Pneumothorax, see Complications
PONV, see Vomiting, nausea and, postoperative
Position, prone, CO_2 tensions during anesthesia, 86:668
Positive inotropic drugs, use after cardiac valve surgery, predictive factors, 86:461
Potency, anesthetic
dose-response
ketorolac, intravenous regional anesthesia, 86:791
midazolam, intramuscular, optimal premedication dose, effects of age and gender, 86:1103
sufentanil, vs epidural fentanyl, dose-response function (letter), 86:918
vecuronium, influence of acute normovolemic hemodilution, 86:861
MAC
desflurane, failure of oleamide to alter, 86:1294
for airway occlusion, effect of vagotomy, halothane-anesthetized cats, 86:191
for airway occlusion, new concept of MAC-airway occlusion response, cats, 86:191
isoflurane, MAC-awake, effect of oral clonidine premedication, 86:410
sevoflurane, effect of adenosine triphosphate, 86:179
sevoflurane, MAC-awake, effect of adenosine triphosphate, 86:179
Power spectrum analysis, see Measurement techniques
Precordial Doppler ultrasound, see Monitoring
Preemptive analgesia, see Anesthetic techniques
Premedication
intramuscular midazolam, optimal dose, effects of age and gender, 86:1103
oral clonidine

effect on hemodynamic responses to pin head-holder application during craniotomy, 86:1001
optimal test dose of epinephrine for epidural injection with lidocaine solution, awake patients, 86:1010
reduction of awakening concentration of isoflurane, 86:410
oral dextromethorphan, reduction of postoperative pain after tonsillectomy, 86:594
oral transmucosal fentanyl citrate (Fentanyl Oralet), pediatric tonsillectomy patients, 86:66
temazepam, effect on hemodynamic responses to pin head-holder application during craniotomy, 86:1001
Preoperative evaluation, history-taking methods, HealthQuiz vs written questionnaire, 86:134
Preservatives, chlorobutanol, effect on contractile force of human atrial trabeculae, 86:40
Propofol, see Anesthetics, intravenous
Prostaglandin E_1, see Pharmacology
Prostate surgery, see Surgery, radical retropubic prostatectomy
Protein binding, bupivacaine and mepivacaine enantiomers, 86:361
Pruritus, see Complications
Pulmonary artery, see Arteries
Pulmonary hypertension, see Complications
Pulse oximetry, see Measurement techniques
Purkinje fibers, see Heart
Pseudothrombocytopenia, see Complications
Pyloric stenosis surgery, see Surgery

Randomized controlled trials, vs observational studies (editorial), 86:225
Receptors
α_2-adrenergic, agonists
clonidine, oral premedication, reduction of awakening concentration of isoflurane, 86:410
clonidine, topical and intravenous, vascular effects, canine pial microcirculation, 86:766
dexmedetomidine, effects in spinal nerve ligation model of neuropathic pain, rats, 86:355
effects on catecholamine activation, vasomotor center, emergence from anesthesia, 86:240
γ-aminobutyric acid
activity, benzodiazepine-sensitive, potentiation by oleamide, 86:1294
gabapentin for neuropathic pain, 86:111
NMDA
antagonists, combined with opioids, analgesic and cognitive effects, 86:1250
antagonists, magnesium sulfate, intrathecal co-infusion with morphine, antinociceptive potentiation and attenuation of tolerance, rats, 86:830
antagonists, MK801, anesthetics and ischemic depolarization in focal ischemia, 86:773
antagonists, potentiation of morphine, 86:825
ketamine delays neuropathic hyperalgesia, rats, 86:557
Recovery
characteristics, desflurane vs sevoflurane, effect of anesthetic duration, volunteers, 86:414
early and late, facilitation, effect of timing of ondansetron administration, ambulatory setting, 86:274
emergence, from anesthesia, catecholamine activation in vasomotor center, effects of α_2 agonists, 86:240
fast-track eligibility, after ambulatory anesthesia, comparison of desflurane, sevoflurane, and propofol, 86:267
from opioid anesthesia, clinical implication of context-sensitive half-times, 86:184
neuromuscular, and mivacurium infusion rate, magnesium-treated parturients, 86:523
profile, patients treated with intrathecal fentanyl with small-dose dilute bupivacaine (letter), 86:917
rate, postoperative, abdominothoracic esophagectomy, 86:228
time
desflurane, "cross-over" from isoflurane toward end of anesthesia, rats breathing from nonrebreathing system, 86:198
isoflurane, "cross-over" to desflurane toward end of anesthesia, rats breathing from nonrebreathing system, 86:198
postoperative, choice of muscle relaxants (letter, reply), 86:1335
Reflexes
skin vasomotor reflex, measured by laser Doppler flowmetry, objective assessment of level of regional anesthesia, 86:336
somatosympathetic, Aδ and C, effect of sevoflurane, dogs, 86:1079
Regional anesthesia and pain management, see Anesthesia
Reintubation, see Intubation, tracheal
Remifentanil, see Anesthetics, intravenous
Research
case-control and cohort studies in anesthesia (letter), 86:1149
informed consent, pediatric anesthesia research, factors that influence parents' decisions, 86:50
Residents, see Education
Respiratory impedance plethysmography, see Measurement techniques
Respiratory inductive plethysmography, see Measurement techniques
Retinal hemorrhage, see Complications
Riluzole, see Pharmacology
Rocuronium, see Neuromuscular relaxants
Ropivacaine, see Anesthetics, local

Sedation
conscious
with chloral hydrate and N_2O, 86:724
with propofol, patient-controlled vs anesthesiologist-controlled, for dental treatment in anxious patients, 86:967
severe trauma patient, safety and efficacy of propofol vs midazolam, 86:1219
Seizures, see Complications
Sensory thresholds, changes after abdominal hysterectomy, anesthetic supplementation with fentanyl vs magnesium or ketamine, 86:95
Sevoflurane, see Anesthetics, volatile
Shoulder surgery, see Surgery
Signal processing, wavelet analysis, embolic heart sound detected by precordial Doppler ultrasound during continuous venous air embolism, dogs, 86:325
Skin vasomotor reflex, see Reflexes
Society for Pediatric Anesthesia, eleventh annual meeting, report, 86:909
Sodium bicarbonate, added to lidocaine
effect on pain during intradermal injection, 86:379
enhancement of depth of epidural blockade, 86:341
Somatosympathetic reflexes, see Reflexes
Spinal anesthesia, see Anesthetic techniques

Spinal cord, accumulation of epidural clonidine, effects of epidural epinephrine, rabbits (letter), 86:1145
Spinal hematoma, see Complications
Splanchnic nerve block, see Anesthetic techniques
Statistics, meta-analysis
acute normovolemic hemodilution and reduction of perioperative allogeneic transfusion, 86:9
comparative effects of postoperative analgesic therapies on pulmonary outcome, 86:598
Sterility
syringes, anesthetic and resuscitative drug, used in obstetric OR, 86:994
thiopental, after preparation, 86:208
Sternocleidomastoid muscle, see Muscle
Steroids, see Pharmacology
Stress hormones, see Hormones
Subclinical portal-systemic encephalopathy, see Complications
Succinylcholine, see Neuromuscular relaxants
Sufentanil
see Analgesics
see Anesthetics, intravenous
Supraclavicular block, see Anesthetic techniques
Supraclavicular nerve block, see Anesthetic techniques
Suprascapular block, see Anesthetic techniques
Surgery
abdominal, lower, postoperative analgesic consumption, effect of epidural verapamil, 86:786
adenotonsillectomy, PONV, antiemetic effectiveness of ondansetron vs dimenhydrinate, children, 86:496
arthroscopic, shoulder, regional anesthesia techniques (letter), 86:217
cardiac
elective, amrinone vs milrinone, multicenter, randomized, blind comparison, 86:683
ethylenediaminetetraacetic acid-dependent pseudothrombocytopenia, 86:1210
IL-6 response, influence of duration of CPB or aortic cross-clamp in absence of blood and/or blood product administration, 86:28
morphine clearance, infants, 86:958
valvular, positive inotropic drug support, predictive factors, 86:461
cardiopulmonary bypass
ϵ-aminocaproic acid plasma levels during (letter), 86:680
CBF velocity and cerebral oxygen extraction, effects of propofol, 86:1201
duration, in absence of blood and/or blood product administration, influence on interleukin-6 response to cardiac surgery, 86:28
gastric mucosal oxygen delivery, decreased, despite constant systemic oxygen delivery, 86:455
hypothermic, cerebral venous oxyhemoglobin desaturation, failure of propofol to ameliorate, 86:926
hypothermic, collapsed anterior chamber (letter), 86:919
influences on hepatic circulation and oxygen metabolism, beagles, 86:1177
ischemic insult, kidney viability, and renal function (editorial), 86:1
normothermic, splanchnic oxygen transport and lactate metabolism, humans, 86:22
pharmacologic electroencephalographic suppression, thiopental vs isoflurane, 86:246
refractory hypotension after, treatment, use of vasopressin, 86:1207
renal protection, patients with preoperative abnormal renal function, 86:3
severe hypercarbia during, complication of CO_2 flooding surgical field, 86:264
systemic inflammatory response, manifestations, 86:28
cerebellar hemangioblastoma excision, patient with von Hippel-Lindau syndrome anesthetic management, 86:673
cesarean section
elective, patients undergoing, intravenous oxytocin (letter, reply), 86:1334
epidural morphine, side effects, droperidol and, 86:532
intrathecal morphine-induced pruritus, subhypnotic doses of propofol do not relieve, 86:310
mivacurium infusion rate and spontaneous neuromuscular recovery, magnesium-treated parturients, 86:523
postoperative pain control, small doses of intrathecal morphine combined with systemic diclofenac, 86:538
sextuplets, successful epidural anesthesia, 86:1236
spinal anesthesia, small-dose hyperbaric vs plain bupivacaine, 86:989
cholecystectomy
laparoscopic, gasless mechanical vs conventional CO_2 pneumoperitoneum methods, 86:153
laparoscopic, incidence of gastroesophageal reflux and tracheal contamination, 86:624
laparoscopic, pneumoperitoneum, -induced left ventricular loading modifications, time course echocardiographic study, 86:482
postoperative analgesia, use of D-myo-inositol-1,2,6-triphosphate, 86:107
coronary artery bypass grafting, minimally invasive, before lung transplantation, anesthetic management, 86:939
craniotomy, pin head-holder application, hemodynamic responses, effect of clonidine premedication, 86:1001
esophagectomy
abdominothoracic, control of postoperative pathophysiology and rehabilitation, multimodal approach, 86:228
cervical, and total laryngectomy, patient with, intra- and postoperative airway management (letter), 86:678
extracorporeal shock wave lithotripsy
intrathecal sufentanil analgesia, patients with aortic stenosis, 86:943
outpatient, general vs epidural anesthesia, 86:1214
gynecological
laparoscopic, diagnostic, incidence of gastroesophageal reflux and tracheal contamination, 86:624
laparoscopic, outpatient, effect of timing of ondansetron administration, 86:274
laparoscopic, pneumoperitoneum as risk factor for endobronchial intubation, 86:301
PONV, prevention, with combination of granisetron and droperidol, 86:613
hand, outpatient, use of selective axillary nerve block, 86:746
hernia repair, efficacy and safety of clonidine/bupivacaine combination, pediatric patients, 86:54

hip, arthroplasty
allogeneic blood requirements, effects of red-cell scavenging, hemodilution, and active warming, 86:387
CO_2 embolism during, infant, 86:299
elective, unsuccessful resuscitation under hypotensive epidural anesthesia, 86:847
hysterectomy, abdominal, postoperative sensory changes and pain, comparison of anesthetic supplementation with fentanyl vs magnesium or ketamine, 86:95
knee
anterior cruciate ligament repair, ambulatory, postoperative pain, multimodal analgesic management, intraarticular morphine, 86:374
arthroplasty, allogeneic blood requirements, effects of red-cell scavenging, hemodilution, and active warming, 86:387
replacement, perioperative clinical pathway, effect on hospital costs, 86:978
laminectomy, lumbar, aortic perforation during, 86:493
laparotomy, management of postoperative pain, alfentanil as adjuvant to epidural bupivacaine, lack of evidence of spinal action, 86:574
laryngectomy, total, and cervical esophagectomy, patient with, intra- and postoperative airway management (letter), 86:678
liver transplantation
fluid requirements, hemodynamic variables, and support during, effects of intraoperative intravenous clonidine, prospective, randomized, study, 86:468
orthotopic, patients undergoing, coagulation, impairment, caused by endogenous heparin-like substances, 86:691
orthotopic, transfusion and coagulation factor requirements, reducing, efficacy of conjugated estrogen, 86:1183
lobectomy, one-lung ventilation, arterial oxygenation and shunt fraction, isoflurane vs sevoflurane, 86:1266
lumbar disk, comparison of PCA with lornoxicam vs morphine, 86:1045
lung transplantation, patient undergoing minimally invasive myocardial revascularization before, anesthetic management, 86:939
myocardial revascularization, minimally invasive, before lung transplantation, anesthetic management, 86:939
noncardiac, morphine clearance, infants, 86:958
open heart, pharmacokinetics and side effects of milrinone, infants and children, 86:283
orchidopexy, laparoscopic, pneumoperitoneum during, hemodynamic effects, assessment by continuous esophageal aortic blood flow echo-Doppler, healthy infants, 86:290
pheochromocytoma, patient with von Hippel-Lindau syndrome anesthetic management, 86:673
pyloric stenosis, awake vs paralyzed tracheal intubation, infants, 86:945
radical retropubic prostatectomy, intravenous lidocaine, effects on postoperative pain, hospital length-of-stay, and return of bowel function, 86:235
shoulder, arthroscopy, outpatient, regional anesthesia techniques (letter), 86:217
thoracic, one-lung ventilation
intrinsic PEEP and flow-volume curves, 86:880
right-sided double-lumen tube vs single-lumen tube with right-sided enclosed bronchial blocker (Univent), 86:696
thoracoscopic, video-assisted, modified endotracheal tube, infants and small children, 86:1212
tonsillectomy, premedication
oral dextromethorphan, reduction of postoperative pain, 86:594
oral transmucosal fentanyl citrate, pediatric patients, 86:66
Sympathetic block, see Anesthetic techniques
Sympathetic nervous system
catecholamines
activation, vasomotor center, emergence from anesthesia, effects of α_2 agonists, 86:240
ephedrine, enantiomers of bupivacaine-lung uptake, 86:812
ephedrine, prophylactic, attenuation of hemodynamic response to propofol, elderly female patients, 86:477
epinephrine, intravenous, simulation of epidural test dose, sevoflurane-anesthetized children, 86:952
clonidine
clonidine/bupivacaine combination, efficacy and safety, caudal blockade for pediatric hernia repair, 86:54
intravenous, intraoperative, effects on fluid requirements, hemodynamic variables, and support during liver transplantation, prospective, randomized, study, 86:468
oral premedication, reduction of awakening concentration of isoflurane, 86:410
sympathetic activity, spontaneous, effect of sevoflurane, dogs, 86:1079
sympathetic blockade and pain from veins, 86:588
Syringes, see Equipment
Systemic inflammatory response, see Inflammation

Tachycardia, see Heart
TEE, see Monitoring, transesophageal echocardiography
Temazepam, see Premedication
Temperature
hypothermia
mild, effect on nitroglycerin-induced vasodilation of pial arteries, cats, 86:546
mild to moderate, arterial to end-tidal CO_2 gradient increases with uncorrected but not with temperature-corrected $Paco_2$ determination, 86:1131
warming, active, effect on allogeneic blood requirements, patients undergoing hip or knee arthroplasty, 86:387
Thermal injury, see Pain
Thiopental, see Anesthetics, intravenous
Thoracic dermatomes, duration of lidocaine spinal anesthesia, effect of varied doses of epinephrine, 86:1018
Thoracic epidural, see Anesthetic techniques
Thoracic surgery, see Surgery
Thoracoscopic surgery, see Surgery
Three-in-one block, see Anesthetic techniques
Thrombocytopenia
see Complications
see Complications, immune thrombocytopenic purpura
see Complications, pseudothrombocytopenia
Thromboelastography, see Monitoring
Thyroid, see Hormones

Tolerance
see Analgesics, remifentanil
see Pain
Tonsillectomy, see Surgery
Torsade de pointes, see Complications
Total intravenous anesthesia, see Anesthetic techniques
Tourniquet
pain, epinephrine and bupivacaine spinal anesthesia, 86:973
release
changes in oxygen consumption and CO_2 elimination, 86:90
time, and analgesic effects of intraarticular morphine (letter), 86:1150
Toxicity
cardiotoxicity, bupivacaine-induced, effects of verapamil and nimodipine, *in vivo* and *in vitro* study, rats, 86:818
hepatic, absence, after four hours of 1.25 MAC sevoflurane anesthesia, volunteers, 86:662
human toxicity study, ethics (letter), 86:445
local anesthetic, treatment, midazolam, flumazenil administered to antagonize, opisthotonos after (letter), 86:677
neurotoxicity, lidocaine, manifestations, graded, irreversible changes in giant axons, crayfish, 86:569
renal, absence, after four hours of 1.25 MAC sevoflurane anesthesia, volunteers, 86:662
sufentanil, ocular application, dogs, 86:138
systemic, levobupivacaine, 86:797
Tracheal cuff, see Equipment, double-lumen tube
Tracheal intubation, see Intubation
Tracheal tube, see Equipment
Transcranial Doppler ultrasonography, see Monitoring
Transesophageal echocardiography, see Monitoring
Transfusion, see Blood
Transient neurologic syndrome, see Complications
Trauma
see Airway, management
see Brain, traumatic injury
see Complications
Tropisetron, see Antiemetics

Ultrasonography
see Monitoring, precordial Doppler ultrasound
see Monitoring, transcranial Doppler ultrasonography
Univent, see Equipment, single-lumen tube
Upper respiratory tract infections, see Complications
Uremic neuropathy, see Complications
Uterus, relaxation, effect of nitroglycerin, enhancement by placental tissue, 86:304

Vascular pain, see Pain
Vasodilation, nitroglycerin-induced, pial arteries, effect of mild hypothermia, cats, 86:546
Vasopressin, see Pharmacology
Vecuronium, see Neuromuscular relaxants
Veins, peripheral, analgesic effect of fentanyl, morphine, meperidine, and lidocaine, comparative study, 86:382
Ventilation
controlled, and endobronchial intubation, management of congenital lobar emphysema, 86:71
depression, sedation with chloral hydrate and N_2O, 86:724
distress, respiratory, induced by airway occlusion, inhibitory influence of pulmonary vagal afferents, halothane-anesthetized cats, 86:398
effects of nonimmobilizers, swine, 86:173
hemidiaphragmatic paresis after supraclavicular block, 86:1239
isocapnic hypoxic ventilatory responses, acute and sustained, during isoflurane anesthesia, 86:403
jet, high-frequency, and LMA, for resection of high tracheal stenosis, 86:907
mechanical, facilitation during fiberoptic intubation, use of jaw support device, 86:432
oxygenation, inhaled prostaglandin E_1, acute lung injury, 86:753
perfusion distribution, effect of dopexamine, anesthetized, paralyzed patients, 86:314
Ventricular dysfunction, see Heart
Verapamil, see Pharmacology
Vision, loss, after lumbar epidural steroid injection, 86:119
Visual analog scale, see Measurement techniques
Vomiting, nausea and, postoperative
adenotonsillectomy, antiemetic effectiveness of ondansetron vs dimenhydrinate, children, 86:496
elective outpatient surgical procedures, antiemetics, ondansetron vs droperidol, efficacy, safety, and patient satisfaction, 86:731
established, treating, use of tropisetron, randomized, double-blind, placebo-controlled study, 86:617
laparoscopy, outpatient, effect of timing of ondansetron administration, 86:274
prevention, combination of granisetron and droperidol, 86:613
von Hippel-Lindau syndrome, see Complications

Warming, see Temperature
Wavelet analysis, see Signal processing

Yohimbine, see Pharmacology